

表 2738 体重減少率(%) %クレアチニン産生速度(%)別(血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
60未満	42	56	113	144	113	73	57	22	7	5	632		632	3.85	1.91
(%)	(6.6)	(8.9)	(17.9)	(22.8)	(17.9)	(11.6)	(9.0)	(3.5)	(1.1)	(0.8)	(100.0)				
60～	30	31	66	115	103	83	62	20	4	3	517	1	518	4.17	1.88
(%)	(5.8)	(6.0)	(12.8)	(22.2)	(19.9)	(16.1)	(12.0)	(3.9)	(0.8)	(0.6)	(100.0)				
70～	37	51	87	194	213	164	102	43	11	5	907		907	4.37	1.80
(%)	(4.1)	(5.6)	(9.6)	(21.4)	(23.5)	(18.1)	(11.2)	(4.7)	(1.2)	(0.6)	(100.0)				
80～	29	58	133	283	338	261	131	69	25	12	1,339		1,339	4.52	1.76
(%)	(2.2)	(4.3)	(9.9)	(21.1)	(25.2)	(19.5)	(9.8)	(5.2)	(1.9)	(0.9)	(100.0)				
90～	30	66	163	368	473	435	225	93	29	10	1,892		1,892	4.67	1.61
(%)	(1.6)	(3.5)	(8.6)	(19.5)	(25.0)	(23.0)	(11.9)	(4.9)	(1.5)	(0.5)	(100.0)				
100～	15	58	183	467	597	532	254	91	31	14	2,242		2,242	4.70	1.49
(%)	(0.7)	(2.6)	(8.2)	(20.8)	(26.6)	(23.7)	(11.3)	(4.1)	(1.4)	(0.6)	(100.0)				
110～	15	50	148	371	605	461	261	96	31	9	2,047		2,047	4.77	1.48
(%)	(0.7)	(2.4)	(7.2)	(18.1)	(29.6)	(22.5)	(12.8)	(4.7)	(1.5)	(0.4)	(100.0)				
120～	11	25	99	301	439	374	178	64	18	12	1,521	1	1,522	4.79	1.44
(%)	(0.7)	(1.6)	(6.5)	(19.8)	(28.9)	(24.6)	(11.7)	(4.2)	(1.2)	(0.8)	(100.0)				
130～	2	12	44	151	246	228	100	33	8	5	829		829	4.87	1.41
(%)	(0.2)	(1.4)	(5.3)	(18.2)	(29.7)	(27.5)	(12.1)	(4.0)	(1.0)	(0.6)	(100.0)				
140～	2	4	36	90	136	118	45	16	9	5	461	7	468	4.78	1.45
(%)	(0.4)	(0.9)	(7.8)	(19.5)	(29.5)	(25.6)	(9.8)	(3.5)	(2.0)	(1.1)	(100.0)				
合計	213	411	1,072	2,484	3,263	2,729	1,415	547	173	80	12,387	9	12,396	4.62	1.61
(%)	(1.7)	(3.3)	(8.7)	(20.1)	(26.3)	(22.0)	(11.4)	(4.4)	(1.4)	(0.6)	(100.0)				
記載なし	11	25	69	103	124	108	45	21	4	3	513	286	799	4.39	1.73
(%)	(2.1)	(4.9)	(13.5)	(20.1)	(24.2)	(21.1)	(8.8)	(4.1)	(0.8)	(0.6)	(100.0)				
総計	224	436	1,141	2,587	3,387	2,837	1,460	568	177	83	12,900	295	13,195	4.61	1.62
(%)	(1.7)	(3.4)	(8.8)	(20.1)	(26.3)	(22.0)	(11.3)	(4.4)	(1.4)	(0.6)	(100.0)				
平均	80.31	89.13	96.01	101.04	105.00	105.75	103.31	101.89	103.22	103.23	102.28	150.73	102.31		
標準偏差	25.79	25.51	26.86	24.26	22.40	21.86	22.98	22.49	24.60	25.77	23.84	39.50	23.89		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査