

表 2742 体重減少率(%) 透析前リン濃度(mg/dL)別(血液透析濾過患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
3未満	41	72	136	220	229	136	88	26	8	4	960	8	968	4.04	1.72
(%)	(4.3)	(7.5)	(14.2)	(22.9)	(23.9)	(14.2)	(9.2)	(2.7)	(0.8)	(0.4)	(100.0)				
3.0～	42	84	173	284	335	230	100	42	11	5	1,306	8	1,314	4.16	1.66
(%)	(3.2)	(6.4)	(13.2)	(21.7)	(25.7)	(17.6)	(7.7)	(3.2)	(0.8)	(0.4)	(100.0)				
3.5～	71	99	289	541	631	458	212	76	22	11	2,410	12	2,422	4.29	1.62
(%)	(2.9)	(4.1)	(12.0)	(22.4)	(26.2)	(19.0)	(8.8)	(3.2)	(0.9)	(0.5)	(100.0)				
4.0～	84	173	437	831	1,117	755	346	125	49	20	3,937	25	3,962	4.39	1.62
(%)	(2.1)	(4.4)	(11.1)	(21.1)	(28.4)	(19.2)	(8.8)	(3.2)	(1.2)	(0.5)	(100.0)				
4.5～	84	158	473	1,091	1,389	1,079	457	167	45	23	4,966	25	4,991	4.49	1.53
(%)	(1.7)	(3.2)	(9.5)	(22.0)	(28.0)	(21.7)	(9.2)	(3.4)	(0.9)	(0.5)	(100.0)				
5.0～	103	174	502	1,142	1,500	1,148	594	194	55	16	5,428	32	5,460	4.52	1.54
(%)	(1.9)	(3.2)	(9.2)	(21.0)	(27.6)	(21.1)	(10.9)	(3.6)	(1.0)	(0.3)	(100.0)				
5.5～	61	114	360	940	1,310	1,060	508	209	54	21	4,637	31	4,668	4.66	1.51
(%)	(1.3)	(2.5)	(7.8)	(20.3)	(28.3)	(22.9)	(11.0)	(4.5)	(1.2)	(0.5)	(100.0)				
6.0～	10	23	59	184	242	189	88	33	7	2	837	7	844	4.61	1.44
(%)	(1.2)	(2.7)	(7.0)	(22.0)	(28.9)	(22.6)	(10.5)	(3.9)	(0.8)	(0.2)	(100.0)				
6.1～	20	59	200	525	807	693	366	136	33	20	2,859	25	2,884	4.81	1.48
(%)	(0.7)	(2.1)	(7.0)	(18.4)	(28.2)	(24.2)	(12.8)	(4.8)	(1.2)	(0.7)	(100.0)				
6.5～	22	44	165	448	668	605	329	127	36	18	2,462	12	2,474	4.87	1.55
(%)	(0.9)	(1.8)	(6.7)	(18.2)	(27.1)	(24.6)	(13.4)	(5.2)	(1.5)	(0.7)	(100.0)				
7.0～	17	27	85	284	441	366	215	92	35	13	1,575	12	1,587	4.92	1.58
(%)	(1.1)	(1.7)	(5.4)	(18.0)	(28.0)	(23.2)	(13.7)	(5.8)	(2.2)	(0.8)	(100.0)				
7.5～	7	20	49	147	251	253	156	58	18	13	972	6	978	5.06	1.57
(%)	(0.7)	(2.1)	(5.0)	(15.1)	(25.8)	(26.0)	(16.0)	(6.0)	(1.9)	(1.3)	(100.0)				
8.0～	8	8	24	89	158	136	80	41	23	7	574	4	578	5.11	1.65
(%)	(1.4)	(1.4)	(4.2)	(15.5)	(27.5)	(23.7)	(13.9)	(7.1)	(4.0)	(1.2)	(100.0)				
8.5～	2	6	17	41	104	88	59	22	17	3	359	5	364	5.18	1.58
(%)	(0.6)	(1.7)	(4.7)	(11.4)	(29.0)	(24.5)	(16.4)	(6.1)	(4.7)	(0.8)	(100.0)				
9.0～	7	7	27	65	134	169	98	56	19	14	596	5	601	5.38	1.71
(%)	(1.2)	(1.2)	(4.5)	(10.9)	(22.5)	(28.4)	(16.4)	(9.4)	(3.2)	(2.3)	(100.0)				
合計	579	1,068	2,996	6,832	9,316	7,365	3,696	1,404	432	190	33,878	217	34,095	4.59	1.58
(%)	(1.7)	(3.2)	(8.8)	(20.2)	(27.5)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
記載なし	8	23	34	61	61	59	29	4	4		283	593	876	4.19	1.65
(%)	(2.8)	(8.1)	(12.0)	(21.6)	(21.6)	(20.8)	(10.2)	(1.4)	(1.4)		(100.0)				
総計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
平均	4.80	4.84	5.00	5.22	5.37	5.53	5.64	5.77	5.89	6.01	5.37	5.50	5.37		
標準偏差	1.40	1.36	1.31	1.31	1.37	1.43	1.51	1.60	1.66	1.87	1.42	1.50	1.42		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査