

表 2743 体重減少率(%) PTH値(pg/mL)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上) PTH測定法:intact-PTH

PTH値	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	25	51	176	313	390	313	129	39	22	5	1,463	8	1,471	4.41	1.57
(%)	(1.7)	(3.5)	(12.0)	(21.4)	(26.7)	(21.4)	(8.8)	(2.7)	(1.5)	(0.3)	(100.0)				
20～	35	84	196	405	504	385	185	76	17	8	1,895	12	1,907	4.44	1.58
(%)	(1.8)	(4.4)	(10.3)	(21.4)	(26.6)	(20.3)	(9.8)	(4.0)	(0.9)	(0.4)	(100.0)				
40～	41	84	181	460	574	461	213	78	24	7	2,123	12	2,135	4.50	1.55
(%)	(1.9)	(4.0)	(8.5)	(21.7)	(27.0)	(21.7)	(10.0)	(3.7)	(1.1)	(0.3)	(100.0)				
60～	44	87	237	440	630	486	185	77	26	11	2,223	7	2,230	4.46	1.58
(%)	(2.0)	(3.9)	(10.7)	(19.8)	(28.3)	(21.9)	(8.3)	(3.5)	(1.2)	(0.5)	(100.0)				
80～	46	80	203	503	694	550	267	99	31	20	2,493	18	2,511	4.60	1.61
(%)	(1.8)	(3.2)	(8.1)	(20.2)	(27.8)	(22.1)	(10.7)	(4.0)	(1.2)	(0.8)	(100.0)				
100～	50	79	230	531	655	501	259	86	36	16	2,443	16	2,459	4.53	1.61
(%)	(2.0)	(3.2)	(9.4)	(21.7)	(26.8)	(20.5)	(10.6)	(3.5)	(1.5)	(0.7)	(100.0)				
120～	50	66	210	465	615	500	229	94	29	8	2,266	15	2,281	4.55	1.60
(%)	(2.2)	(2.9)	(9.3)	(20.5)	(27.1)	(22.1)	(10.1)	(4.1)	(1.3)	(0.4)	(100.0)				
140～	36	47	148	416	573	461	219	90	27	6	2,023	12	2,035	4.65	1.50
(%)	(1.8)	(2.3)	(7.3)	(20.6)	(28.3)	(22.8)	(10.8)	(4.4)	(1.3)	(0.3)	(100.0)				
160～	29	58	169	372	527	398	207	83	17	16	1,876	16	1,892	4.62	1.59
(%)	(1.5)	(3.1)	(9.0)	(19.8)	(28.1)	(21.2)	(11.0)	(4.4)	(0.9)	(0.9)	(100.0)				
180～	63	130	347	845	1,207	980	496	204	48	32	4,352	28	4,380	4.67	1.60
(%)	(1.4)	(3.0)	(8.0)	(19.4)	(27.7)	(22.5)	(11.4)	(4.7)	(1.1)	(0.7)	(100.0)				
240～	1	3	6	8	24	21	4	2		1	70		70	4.61	1.51
(%)	(1.4)	(4.3)	(8.6)	(11.4)	(34.3)	(30.0)	(5.7)	(2.9)		(1.4)	(100.0)				
241～	30	47	193	508	743	563	293	124	45	9	2,555	20	2,575	4.73	1.51
(%)	(1.2)	(1.8)	(7.6)	(19.9)	(29.1)	(22.0)	(11.5)	(4.9)	(1.8)	(0.4)	(100.0)				
300～	19	38	119	263	406	344	198	66	20	8	1,481	8	1,489	4.74	1.54
(%)	(1.3)	(2.6)	(8.0)	(17.8)	(27.4)	(23.2)	(13.4)	(4.5)	(1.4)	(0.5)	(100.0)				
360～	10	23	79	157	248	224	112	41	22	5	921	4	925	4.76	1.58
(%)	(1.1)	(2.5)	(8.6)	(17.0)	(26.9)	(24.3)	(12.2)	(4.5)	(2.4)	(0.5)	(100.0)				
420～	15	14	44	84	124	143	59	25	8	5	521	3	524	4.70	1.69
(%)	(2.9)	(2.7)	(8.4)	(16.1)	(23.8)	(27.4)	(11.3)	(4.8)	(1.5)	(1.0)	(100.0)				
480～	2	6	10	17	31	30	23	3	2	1	125		125	4.83	1.72
(%)	(1.6)	(4.8)	(8.0)	(13.6)	(24.8)	(24.0)	(18.4)	(2.4)	(1.6)	(0.8)	(100.0)				
500～	1	3	10	35	48	53	38	12	2		202	2	204	5.01	1.42
(%)	(0.5)	(1.5)	(5.0)	(17.3)	(23.8)	(26.2)	(18.8)	(5.9)	(1.0)		(100.0)				
540～	1	4	18	41	60	46	36	13	6	5	230	2	232	4.98	1.68
(%)	(0.4)	(1.7)	(7.8)	(17.8)	(26.1)	(20.0)	(15.7)	(5.7)	(2.6)	(2.2)	(100.0)				
600～	7	4	19	65	110	83	59	31	9	4	391		391	5.06	1.64
(%)	(1.8)	(1.0)	(4.9)	(16.6)	(28.1)	(21.2)	(15.1)	(7.9)	(2.3)	(1.0)	(100.0)				
800～	5	9	25	51	74	75	59	21	4	7	330	4	334	5.00	1.78
(%)	(1.5)	(2.7)	(7.6)	(15.5)	(22.4)	(22.7)	(17.9)	(6.4)	(1.2)	(2.1)	(100.0)				
合計	510	917	2,620	5,979	8,237	6,617	3,270	1,264	395	174	29,983	187	30,170	4.61	1.58
(%)	(1.7)	(3.1)	(8.7)	(19.9)	(27.5)	(22.1)	(10.9)	(4.2)	(1.3)	(0.6)	(100.0)				
記載なし	13	32	48	108	142	83	30	9	9	2	476	100	576	4.22	1.69
(%)	(2.7)	(6.7)	(10.1)	(22.7)	(29.8)	(17.4)	(6.3)	(1.9)	(1.9)	(0.4)	(100.0)				
総計	523	949	2,668	6,087	8,379	6,700	3,300	1,273	404	176	30,459	287	30,746	4.60	1.59
(%)	(1.7)	(3.1)	(8.8)	(20.0)	(27.5)	(22.0)	(10.8)	(4.2)	(1.3)	(0.6)	(100.0)				
平均	167.73	155.20	165.27	169.97	175.71	182.23	200.78	203.32	196.29	224.76	178.78	180.86	178.80		
標準偏差	213.70	152.61	180.01	176.16	155.36	168.27	197.87	202.21	169.92	230.34	173.81	156.99	173.71		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy
2014年調査