

表 2745 体重減少率(%) 透析前ヘモグロビン濃度(g/dL)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

透析前ヘモグロビン濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
5未満		1		5	9	4	4	1			24		24	4.73	1.28
(%)		(4.2)		(20.8)	(37.5)	(16.7)	(16.7)	(4.2)			(100.0)				
5～	2	1	5	5	5	9	1				28		28	3.90	1.66
(%)	(7.1)	(3.6)	(17.9)	(17.9)	(17.9)	(32.1)	(3.6)				(100.0)				
6～	4	4	10	16	20	19	14	4	2		93	1	94	4.53	1.77
(%)	(4.3)	(4.3)	(10.8)	(17.2)	(21.5)	(20.4)	(15.1)	(4.3)	(2.2)		(100.0)				
7～	11	15	42	53	84	71	50	16	8	2	352	9	361	4.61	1.80
(%)	(3.1)	(4.3)	(11.9)	(15.1)	(23.9)	(20.2)	(14.2)	(4.5)	(2.3)	(0.6)	(100.0)				
8～	23	49	121	246	353	283	161	67	27	11	1,341	22	1,363	4.68	1.72
(%)	(1.7)	(3.7)	(9.0)	(18.3)	(26.3)	(21.1)	(12.0)	(5.0)	(2.0)	(0.8)	(100.0)				
9～	88	186	480	980	1,303	1,060	604	230	87	27	5,045	38	5,083	4.61	1.64
(%)	(1.7)	(3.7)	(9.5)	(19.4)	(25.8)	(21.0)	(12.0)	(4.6)	(1.7)	(0.5)	(100.0)				
10～	187	345	941	2,323	3,134	2,455	1,227	512	136	69	11,329	60	11,389	4.62	1.59
(%)	(1.7)	(3.0)	(8.3)	(20.5)	(27.7)	(21.7)	(10.8)	(4.5)	(1.2)	(0.6)	(100.0)				
11～	165	289	895	2,054	2,868	2,269	1,105	378	107	56	10,186	48	10,234	4.59	1.55
(%)	(1.6)	(2.8)	(8.8)	(20.2)	(28.2)	(22.3)	(10.8)	(3.7)	(1.1)	(0.5)	(100.0)				
12～	67	114	316	780	1,105	817	391	147	55	20	3,812	25	3,837	4.57	1.56
(%)	(1.8)	(3.0)	(8.3)	(20.5)	(29.0)	(21.4)	(10.3)	(3.9)	(1.4)	(0.5)	(100.0)				
13～	16	40	105	228	273	262	85	38	8	1	1,056	9	1,065	4.47	1.49
(%)	(1.5)	(3.8)	(9.9)	(21.6)	(25.9)	(24.8)	(8.0)	(3.6)	(0.8)	(0.1)	(100.0)				
14～	7	18	56	120	126	90	44	9	2	1	473	5	478	4.30	1.45
(%)	(1.5)	(3.8)	(11.8)	(25.4)	(26.6)	(19.0)	(9.3)	(1.9)	(0.4)	(0.2)	(100.0)				
合計	570	1,062	2,971	6,810	9,280	7,339	3,686	1,402	432	187	33,739	217	33,956	4.60	1.58
(%)	(1.7)	(3.1)	(8.8)	(20.2)	(27.5)	(21.8)	(10.9)	(4.2)	(1.3)	(0.6)	(100.0)				
記載なし	17	29	59	83	97	85	39	6	4	3	422	593	1,015	4.18	1.72
(%)	(4.0)	(6.9)	(14.0)	(19.7)	(23.0)	(20.1)	(9.2)	(1.4)	(0.9)	(0.7)	(100.0)				
総計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
平均	10.78	10.79	10.85	10.90	10.88	10.87	10.77	10.72	10.64	10.70	10.85	10.61	10.85		
標準偏差	1.36	1.37	1.32	1.26	1.23	1.24	1.25	1.21	1.24	1.12	1.26	1.52	1.26		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査