

表 2746 体重減少率(%) 総コレステロール濃度(mg/dL)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

総コレステロール濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
100未満	20	31	104	188	197	174	82	31	15	6	848	6	854	4.46	1.67
(%)	(2.4)	(3.7)	(12.3)	(22.2)	(23.2)	(20.5)	(9.7)	(3.7)	(1.8)	(0.7)	(100.0)				
100～	41	88	241	549	739	561	244	97	21	11	2,592	9	2,601	4.50	1.53
(%)	(1.6)	(3.4)	(9.3)	(21.2)	(28.5)	(21.6)	(9.4)	(3.7)	(0.8)	(0.4)	(100.0)				
120～	83	139	466	960	1,406	1,083	535	190	67	26	4,955	28	4,983	4.59	1.59
(%)	(1.7)	(2.8)	(9.4)	(19.4)	(28.4)	(21.9)	(10.8)	(3.8)	(1.4)	(0.5)	(100.0)				
140～	93	187	530	1,222	1,634	1,324	665	233	94	44	6,026	32	6,058	4.62	1.59
(%)	(1.5)	(3.1)	(8.8)	(20.3)	(27.1)	(22.0)	(11.0)	(3.9)	(1.6)	(0.7)	(100.0)				
160～	96	163	466	1,117	1,448	1,087	567	226	60	30	5,260	22	5,282	4.58	1.58
(%)	(1.8)	(3.1)	(8.9)	(21.2)	(27.5)	(20.7)	(10.8)	(4.3)	(1.1)	(0.6)	(100.0)				
180～	64	94	305	694	893	807	363	150	44	12	3,426	15	3,441	4.60	1.58
(%)	(1.9)	(2.7)	(8.9)	(20.3)	(26.1)	(23.6)	(10.6)	(4.4)	(1.3)	(0.4)	(100.0)				
200～	38	82	161	414	554	430	216	79	22	7	2,003	4	2,007	4.53	1.57
(%)	(1.9)	(4.1)	(8.0)	(20.7)	(27.7)	(21.5)	(10.8)	(3.9)	(1.1)	(0.3)	(100.0)				
220～	20	27	75	161	255	169	91	35	10	5	848	2	850	4.54	1.61
(%)	(2.4)	(3.2)	(8.8)	(19.0)	(30.1)	(19.9)	(10.7)	(4.1)	(1.2)	(0.6)	(100.0)				
240～	10	15	37	64	93	63	34	15	8		339		339	4.48	1.69
(%)	(2.9)	(4.4)	(10.9)	(18.9)	(27.4)	(18.6)	(10.0)	(4.4)	(2.4)		(100.0)				
260～	3	5	17	32	48	46	24	6	1	3	185	1	186	4.70	1.58
(%)	(1.6)	(2.7)	(9.2)	(17.3)	(25.9)	(24.9)	(13.0)	(3.2)	(0.5)	(1.6)	(100.0)				
合計	468	831	2,402	5,401	7,267	5,744	2,821	1,062	342	144	26,482	119	26,601	4.58	1.58
(%)	(1.8)	(3.1)	(9.1)	(20.4)	(27.4)	(21.7)	(10.7)	(4.0)	(1.3)	(0.5)	(100.0)				
記載なし	119	260	628	1,492	2,110	1,680	904	346	94	46	7,679	691	8,370	4.65	1.59
(%)	(1.5)	(3.4)	(8.2)	(19.4)	(27.5)	(21.9)	(11.8)	(4.5)	(1.2)	(0.6)	(100.0)				
総計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
平均	161.08	159.45	156.74	157.92	158.25	158.23	159.17	159.42	158.19	157.18	158.27	151.94	158.24		
標準偏差	37.56	37.03	36.49	35.39	35.43	35.61	35.41	35.03	34.92	37.87	35.64	32.87	35.63		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査