

表 2748 体重減少率(%) Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

Non-HDL-C濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
50未満	6	16	48	82	103	101	60	29	12	4	461	2	463	4.79	1.71
(%)	(1.3)	(3.5)	(10.4)	(17.8)	(22.3)	(21.9)	(13.0)	(6.3)	(2.6)	(0.9)	(100.0)				
50～	34	74	157	386	550	496	272	112	34	11	2,126	9	2,135	4.74	1.61
(%)	(1.6)	(3.5)	(7.4)	(18.2)	(25.9)	(23.3)	(12.8)	(5.3)	(1.6)	(0.5)	(100.0)				
70～	79	118	388	893	1,198	1,017	522	186	76	31	4,508	25	4,533	4.65	1.60
(%)	(1.8)	(2.6)	(8.6)	(19.8)	(26.6)	(22.6)	(11.6)	(4.1)	(1.7)	(0.7)	(100.0)				
90～	79	167	523	1,113	1,520	1,175	593	217	70	33	5,490	29	5,519	4.59	1.60
(%)	(1.4)	(3.0)	(9.5)	(20.3)	(27.7)	(21.4)	(10.8)	(4.0)	(1.3)	(0.6)	(100.0)				
110～	90	138	420	981	1,264	967	447	166	51	22	4,546	14	4,560	4.51	1.55
(%)	(2.0)	(3.0)	(9.2)	(21.6)	(27.8)	(21.3)	(9.8)	(3.7)	(1.1)	(0.5)	(100.0)				
130～	59	96	256	604	787	583	244	91	25	9	2,754	12	2,766	4.44	1.53
(%)	(2.1)	(3.5)	(9.3)	(21.9)	(28.6)	(21.2)	(8.9)	(3.3)	(0.9)	(0.3)	(100.0)				
150～	24	54	148	320	384	295	128	46	7	3	1,409	4	1,413	4.39	1.50
(%)	(1.7)	(3.8)	(10.5)	(22.7)	(27.3)	(20.9)	(9.1)	(3.3)	(0.5)	(0.2)	(100.0)				
170～	16	26	73	129	182	97	45	19	6	1	594	1	595	4.24	1.57
(%)	(2.7)	(4.4)	(12.3)	(21.7)	(30.6)	(16.3)	(7.6)	(3.2)	(1.0)	(0.2)	(100.0)				
190～	5	5	13	55	60	47	26	3		1	215	1	216	4.48	1.40
(%)	(2.3)	(2.3)	(6.0)	(25.6)	(27.9)	(21.9)	(12.1)	(1.4)		(0.5)	(100.0)				
210～	2	5	14	23	35	22	12	1		1	115		115	4.29	1.48
(%)	(1.7)	(4.3)	(12.2)	(20.0)	(30.4)	(19.1)	(10.4)	(0.9)		(0.9)	(100.0)				
合計	394	699	2,040	4,586	6,083	4,800	2,349	870	281	116	22,218	97	22,315	4.56	1.58
(%)	(1.8)	(3.1)	(9.2)	(20.6)	(27.4)	(21.6)	(10.6)	(3.9)	(1.3)	(0.5)	(100.0)				
記載なし	193	392	990	2,307	3,294	2,624	1,376	538	155	74	11,943	713	12,656	4.65	1.60
(%)	(1.6)	(3.3)	(8.3)	(19.3)	(27.6)	(22.0)	(11.5)	(4.5)	(1.3)	(0.6)	(100.0)				
総計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
平均	110.40	109.28	108.99	108.88	108.10	105.71	103.47	101.93	97.01	99.75	106.99	101.39	106.97		
標準偏差	33.75	34.64	33.79	33.13	33.09	32.95	33.07	32.01	29.83	32.71	33.19	30.33	33.18		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査