

表 2755 Body Mass Index 併用の状況別（透析患者全体）

併用の状況	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	1,558	9,113	29,960	53,189	58,692	45,119	27,029	14,353	7,401	6,385	252,799	47,995	300,794	21.47	3.65
(%)	(0.6)	(3.6)	(11.9)	(21.0)	(23.2)	(17.8)	(10.7)	(5.7)	(2.9)	(2.5)	(100.0)				
PD単独	25	65	193	480	867	945	687	433	224	161	4,080	3,108	7,188	23.07	3.64
(%)	(0.6)	(1.6)	(4.7)	(11.8)	(21.3)	(23.2)	(16.8)	(10.6)	(5.5)	(3.9)	(100.0)				
HD(F)単独.PD力テ あり	1	4	29	30	68	30	13	8	10	4	197	72	269	21.33	3.50
(%)	(0.5)	(2.0)	(14.7)	(15.2)	(34.5)	(15.2)	(6.6)	(4.1)	(5.1)	(2.0)	(100.0)				
PD.HD併用週1 HD(F)	1	4	31	136	232	291	195	110	68	50	1,118	426	1,544	23.37	3.39
(%)	(0.1)	(0.4)	(2.8)	(12.2)	(20.8)	(26.0)	(17.4)	(9.8)	(6.1)	(4.5)	(100.0)				
PD.HD併用週2 HD(F)	1	2	6	10	32	18	24	16	9	9	127	50	177	23.61	4.00
(%)	(0.8)	(1.6)	(4.7)	(7.9)	(25.2)	(14.2)	(18.9)	(12.6)	(7.1)	(7.1)	(100.0)				
PD.HD併用週3 HD(F)		1	1	5	7	6	6	1	2		29	11	40	22.14	3.34
(%)		(3.4)	(3.4)	(17.2)	(24.1)	(20.7)	(20.7)	(3.4)	(6.9)		(100.0)				
PD.HD併用週4 HD(F)					1						1		1	21.96	
(%)					(100.0)						(100.0)				
PD.HD併用その他		1	1	5	17	11	11	6	2	1	55	18	73	22.88	3.33
(%)		(1.8)	(1.8)	(9.1)	(30.9)	(20.0)	(20.0)	(10.9)	(3.6)	(1.8)	(100.0)				
合計	1,586	9,190	30,221	53,855	59,916	46,420	27,965	14,927	7,716	6,610	258,406	51,680	310,086	21.51	3.66
(%)	(0.6)	(3.6)	(11.7)	(20.8)	(23.2)	(18.0)	(10.8)	(5.8)	(3.0)	(2.6)	(100.0)				
不明			2	2	4	1	1	2		1	13	9	22	21.98	4.28
(%)			(15.4)	(15.4)	(30.8)	(7.7)	(7.7)	(15.4)		(7.7)	(100.0)				
記載なし (%)															
総計	1,586	9,190	30,223	53,857	59,920	46,421	27,966	14,929	7,716	6,611	258,419	51,689	310,108	21.51	3.66
(%)	(0.6)	(3.6)	(11.7)	(20.8)	(23.2)	(18.0)	(10.8)	(5.8)	(3.0)	(2.6)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査