

表 2761 Body Mass Index 週透析回数(回/週)別（透析患者全体）

週透析回数	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
1回	11	25	89	238	319	368	240	131	70	58	1,549	291	1,840	22.65	3.66
(%)	(0.7)	(1.6)	(5.7)	(15.4)	(20.6)	(23.8)	(15.5)	(8.5)	(4.5)	(3.7)	(100.0)				
2回	67	265	746	1,166	1,316	863	520	229	113	95	5,380	812	6,192	20.94	3.54
(%)	(1.2)	(4.9)	(13.9)	(21.7)	(24.5)	(16.0)	(9.7)	(4.3)	(2.1)	(1.8)	(100.0)				
3回	1,465	8,715	28,754	51,240	56,578	43,600	26,101	13,929	7,185	6,188	243,755	26,884	270,639	21.48	3.65
(%)	(0.6)	(3.6)	(11.8)	(21.0)	(23.2)	(17.9)	(10.7)	(5.7)	(2.9)	(2.5)	(100.0)				
4回	2	22	94	149	186	167	106	67	40	35	868	126	994	22.22	3.87
(%)	(0.2)	(2.5)	(10.8)	(17.2)	(21.4)	(19.2)	(12.2)	(7.7)	(4.6)	(4.0)	(100.0)				
5回			11	11	17	8	9	6	1	1	64	25	89	21.75	3.60
(%)			(17.2)	(17.2)	(26.6)	(12.5)	(14.1)	(9.4)	(1.6)	(1.6)	(100.0)				
6回			8	18	11	11	9	2	2	1	62	19	81	21.57	3.32
(%)			(12.9)	(29.0)	(17.7)	(17.7)	(14.5)	(3.2)	(3.2)	(1.6)	(100.0)				
7回			1	4	6	1	6	1	1	2	22	8	30	23.13	3.76
(%)			(4.5)	(18.2)	(27.3)	(4.5)	(27.3)	(4.5)	(4.5)	(9.1)	(100.0)				
合計	1,545	9,027	29,703	52,826	58,433	45,018	26,991	14,365	7,412	6,380	251,700	28,165	279,865	21.48	3.65
(%)	(0.6)	(3.6)	(11.8)	(21.0)	(23.2)	(17.9)	(10.7)	(5.7)	(2.9)	(2.5)	(100.0)				
記載なし	41	163	520	1,031	1,487	1,403	975	564	304	231	6,719	23,524	30,243	22.43	3.74
(%)	(0.6)	(2.4)	(7.7)	(15.3)	(22.1)	(20.9)	(14.5)	(8.4)	(4.5)	(3.4)	(100.0)				
総計	1,586	9,190	30,223	53,857	59,920	46,421	27,966	14,929	7,716	6,611	258,419	51,689	310,108	21.51	3.66
(%)	(0.6)	(3.6)	(11.7)	(20.8)	(23.2)	(18.0)	(10.8)	(5.8)	(3.0)	(2.6)	(100.0)				
平均	2.94	2.97	2.97	2.97	2.97	2.97	2.97	2.97	2.97	2.97	2.97	2.96	2.97		
標準偏差	0.26	0.20	0.21	0.22	0.23	0.24	0.26	0.25	0.25	0.25	0.23	0.30	0.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査