

表 2766 Body Mass Index 透析前リン濃度(mg/dL)別 (透析患者全体)

透析前リン濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
3未満	204	851	1,855	2,243	1,976	1,283	676	295	137	100	9,620	1,266	10,886	20.02	3.50
(%)	(2.1)	(8.8)	(19.3)	(23.3)	(20.5)	(13.3)	(7.0)	(3.1)	(1.4)	(1.0)	(100.0)				
3.0～	142	677	1,841	2,884	2,785	1,957	1,076	488	239	168	12,257	1,325	13,582	20.72	3.48
(%)	(1.2)	(5.5)	(15.0)	(23.5)	(22.7)	(16.0)	(8.8)	(4.0)	(1.9)	(1.4)	(100.0)				
3.5～	172	1,010	3,012	4,853	5,070	3,638	2,040	1,026	447	361	21,629	2,053	23,682	20.99	3.50
(%)	(0.8)	(4.7)	(13.9)	(22.4)	(23.4)	(16.8)	(9.4)	(4.7)	(2.1)	(1.7)	(100.0)				
4.0～	196	1,275	4,154	7,192	7,765	5,692	3,172	1,616	770	588	32,420	2,862	35,282	21.19	3.49
(%)	(0.6)	(3.9)	(12.8)	(22.2)	(24.0)	(17.6)	(9.8)	(5.0)	(2.4)	(1.8)	(100.0)				
4.5～	201	1,328	4,484	8,362	9,273	7,008	4,157	2,059	1,071	829	38,772	3,287	42,059	21.42	3.54
(%)	(0.5)	(3.4)	(11.6)	(21.6)	(23.9)	(18.1)	(10.7)	(5.3)	(2.8)	(2.1)	(100.0)				
5.0～	179	1,250	4,426	8,173	9,410	7,240	4,453	2,270	1,126	890	39,417	3,301	42,718	21.55	3.55
(%)	(0.5)	(3.2)	(11.2)	(20.7)	(23.9)	(18.4)	(11.3)	(5.8)	(2.9)	(2.3)	(100.0)				
5.5～	170	922	3,660	6,950	7,906	6,347	3,914	2,067	1,059	920	33,915	2,790	36,705	21.70	3.63
(%)	(0.5)	(2.7)	(10.8)	(20.5)	(23.3)	(18.7)	(11.5)	(6.1)	(3.1)	(2.7)	(100.0)				
6.0～	27	170	631	1,094	1,362	1,170	631	388	171	150	5,794	467	6,261	21.74	3.62
(%)	(0.5)	(2.9)	(10.9)	(18.9)	(23.5)	(20.2)	(10.9)	(6.7)	(3.0)	(2.6)	(100.0)				
6.1～	82	525	2,004	3,812	4,331	3,755	2,246	1,309	695	615	19,374	1,735	21,109	21.88	3.71
(%)	(0.4)	(2.7)	(10.3)	(19.7)	(22.4)	(19.4)	(11.6)	(6.8)	(3.6)	(3.2)	(100.0)				
6.5～	81	426	1,577	3,178	3,899	3,115	2,039	1,200	689	589	16,793	1,485	18,278	22.05	3.77
(%)	(0.5)	(2.5)	(9.4)	(18.9)	(23.2)	(18.5)	(12.1)	(7.1)	(4.1)	(3.5)	(100.0)				
7.0～	29	264	921	1,948	2,278	1,906	1,310	780	432	426	10,294	983	11,277	22.22	3.84
(%)	(0.3)	(2.6)	(8.9)	(18.9)	(22.1)	(18.5)	(12.7)	(7.6)	(4.2)	(4.1)	(100.0)				
7.5～	33	171	558	1,086	1,367	1,175	828	482	285	315	6,300	558	6,858	22.36	3.99
(%)	(0.5)	(2.7)	(8.9)	(17.2)	(21.7)	(18.7)	(13.1)	(7.7)	(4.5)	(5.0)	(100.0)				
8.0～	12	83	317	627	781	689	471	317	173	224	3,694	359	4,053	22.61	4.06
(%)	(0.3)	(2.2)	(8.6)	(17.0)	(21.1)	(18.7)	(12.8)	(8.6)	(4.7)	(6.1)	(100.0)				
8.5～	6	52	202	396	465	444	304	193	106	137	2,305	267	2,572	22.62	4.10
(%)	(0.3)	(2.3)	(8.8)	(17.2)	(20.2)	(19.3)	(13.2)	(8.4)	(4.6)	(5.9)	(100.0)				
9.0～	25	96	295	578	768	624	452	317	253	251	3,659	418	4,077	22.80	4.31
(%)	(0.7)	(2.6)	(8.1)	(15.8)	(21.0)	(17.1)	(12.4)	(8.7)	(6.9)	(6.9)	(100.0)				
合計	1,559	9,100	29,937	53,376	59,436	46,043	27,769	14,807	7,653	6,563	256,243	23,156	279,399	21.51	3.66
(%)	(0.6)	(3.6)	(11.7)	(20.8)	(23.2)	(18.0)	(10.8)	(5.8)	(3.0)	(2.6)	(100.0)				
記載なし	27	90	286	481	484	378	197	122	63	48	2,176	28,533	30,709	21.20	3.70
(%)	(1.2)	(4.1)	(13.1)	(22.1)	(22.2)	(17.4)	(9.1)	(5.6)	(2.9)	(2.2)	(100.0)				
総計	1,586	9,190	30,223	53,857	59,920	46,421	27,966	14,929	7,716	6,611	258,419	51,689	310,108	21.51	3.66
(%)	(0.6)	(3.6)	(11.7)	(20.8)	(23.2)	(18.0)	(10.8)	(5.8)	(3.0)	(2.6)	(100.0)				
平均	4.67	4.79	4.97	5.11	5.20	5.29	5.38	5.50	5.63	5.80	5.22	5.17	5.21		
標準偏差	1.62	1.50	1.41	1.38	1.37	1.38	1.40	1.44	1.53	1.57	1.42	1.53	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査