

表 2768 Body Mass Index PTH値(pg/mL)別 (透析患者全体) PTH測定法:whole-PTH

PTH値	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
20未満	27	104	259	408	361	293	161	85	43	34	1,775	119	1,894	20.89	3.68
(%)	(1.5)	(5.9)	(14.6)	(23.0)	(20.3)	(16.5)	(9.1)	(4.8)	(2.4)	(1.9)	(100.0)				
20～	22	95	294	463	520	352	264	104	47	38	2,199	132	2,331	21.16	3.54
(%)	(1.0)	(4.3)	(13.4)	(21.1)	(23.6)	(16.0)	(12.0)	(4.7)	(2.1)	(1.7)	(100.0)				
35～	22	68	242	424	401	325	224	101	47	38	1,892	151	2,043	21.28	3.61
(%)	(1.2)	(3.6)	(12.8)	(22.4)	(21.2)	(17.2)	(11.8)	(5.3)	(2.5)	(2.0)	(100.0)				
50～	14	95	325	629	697	568	326	163	82	55	2,954	199	3,153	21.47	3.49
(%)	(0.5)	(3.2)	(11.0)	(21.3)	(23.6)	(19.2)	(11.0)	(5.5)	(2.8)	(1.9)	(100.0)				
75～	7	93	265	505	617	467	304	156	77	71	2,562	186	2,748	21.70	3.62
(%)	(0.3)	(3.6)	(10.3)	(19.7)	(24.1)	(18.2)	(11.9)	(6.1)	(3.0)	(2.8)	(100.0)				
100～	7	61	193	466	503	440	233	115	68	65	2,151	147	2,298	21.76	3.60
(%)	(0.3)	(2.8)	(9.0)	(21.7)	(23.4)	(20.5)	(10.8)	(5.3)	(3.2)	(3.0)	(100.0)				
125～	3	38	141	262	330	309	172	83	39	41	1,418	121	1,539	21.88	3.57
(%)	(0.2)	(2.7)	(9.9)	(18.5)	(23.3)	(21.8)	(12.1)	(5.9)	(2.8)	(2.9)	(100.0)				
150～			7	6	10	11	6	5	1	1	47	1	48	22.02	3.46
(%)			(14.9)	(12.8)	(21.3)	(23.4)	(12.8)	(10.6)	(2.1)	(2.1)	(100.0)				
151～	5	24	105	223	294	216	151	73	43	32	1,166	97	1,263	21.95	3.59
(%)	(0.4)	(2.1)	(9.0)	(19.1)	(25.2)	(18.5)	(13.0)	(6.3)	(3.7)	(2.7)	(100.0)				
180～	4	17	86	164	179	160	97	53	31	24	815	60	875	21.92	3.62
(%)	(0.5)	(2.1)	(10.6)	(20.1)	(22.0)	(19.6)	(11.9)	(6.5)	(3.8)	(2.9)	(100.0)				
210～	2	15	56	105	116	120	74	38	23	32	581	47	628	22.27	3.97
(%)	(0.3)	(2.6)	(9.6)	(18.1)	(20.0)	(20.7)	(12.7)	(6.5)	(4.0)	(5.5)	(100.0)				
240～	1	17	57	127	138	95	81	37	18	26	597	77	674	21.98	3.78
(%)	(0.2)	(2.8)	(9.5)	(21.3)	(23.1)	(15.9)	(13.6)	(6.2)	(3.0)	(4.4)	(100.0)				
300～		4	32	35	50	45	26	33	20	10	255	29	284	22.78	4.14
(%)		(1.6)	(12.5)	(13.7)	(19.6)	(17.6)	(10.2)	(12.9)	(7.8)	(3.9)	(100.0)				
350～	1	6	14	27	35	30	17	12	4	7	153	26	179	21.98	3.86
(%)	(0.7)	(3.9)	(9.2)	(17.6)	(22.9)	(19.6)	(11.1)	(7.8)	(2.6)	(4.6)	(100.0)				
400～	1	5	25	37	32	38	18	13	9	6	184	18	202	21.82	3.90
(%)	(0.5)	(2.7)	(13.6)	(20.1)	(17.4)	(20.7)	(9.8)	(7.1)	(4.9)	(3.3)	(100.0)				
500～	1	9	35	50	49	33	27	12	8	4	228	19	247	21.23	3.62
(%)	(0.4)	(3.9)	(15.4)	(21.9)	(21.5)	(14.5)	(11.8)	(5.3)	(3.5)	(1.8)	(100.0)				
合計	117	651	2,136	3,931	4,332	3,502	2,181	1,083	560	484	18,977	1,429	20,406	21.57	3.63
(%)	(0.6)	(3.4)	(11.3)	(20.7)	(22.8)	(18.5)	(11.5)	(5.7)	(3.0)	(2.6)	(100.0)				
記載なし	5	23	60	77	87	61	24	11	9	5	362	295	657	20.52	3.47
(%)	(1.4)	(6.4)	(16.6)	(21.3)	(24.0)	(16.9)	(6.6)	(3.0)	(2.5)	(1.4)	(100.0)				
総計	122	674	2,196	4,008	4,419	3,563	2,205	1,094	569	489	19,339	1,724	21,063	21.55	3.63
(%)	(0.6)	(3.5)	(11.4)	(20.7)	(22.9)	(18.4)	(11.4)	(5.7)	(2.9)	(2.5)	(100.0)				
平均	72.54	93.53	105.17	103.53	105.88	108.15	108.54	118.07	122.31	126.20	107.11	119.95	108.01		
標準偏差	105.02	116.79	139.25	119.11	106.89	106.05	100.13	123.75	110.83	105.90	114.24	110.38	114.01		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査