

表 2779 Body Mass Index 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	471	2,714	9,027	15,016	15,065	10,134	5,379	2,526	1,134	955	62,421	6,156	68,577	20.84	3.40
(%)	(0.8)	(4.3)	(14.5)	(24.1)	(24.1)	(16.2)	(8.6)	(4.0)	(1.8)	(1.5)	(100.0)				
慢性腎盂腎炎	32	135	323	440	426	299	147	75	29	26	1,932	233	2,165	20.44	3.53
(%)	(1.7)	(7.0)	(16.7)	(22.8)	(22.0)	(15.5)	(7.6)	(3.9)	(1.5)	(1.3)	(100.0)				
急速進行性糸球体腎炎	15	119	274	454	386	222	108	44	24	10	1,656	178	1,834	20.13	3.17
(%)	(0.9)	(7.2)	(16.5)	(27.4)	(23.3)	(13.4)	(6.5)	(2.7)	(1.4)	(0.6)	(100.0)				
妊娠腎/妊娠中毒症	10	68	161	243	213	145	81	43	23	22	1,009	55	1,064	20.68	3.69
(%)	(1.0)	(6.7)	(16.0)	(24.1)	(21.1)	(14.4)	(8.0)	(4.3)	(2.3)	(2.2)	(100.0)				
その他の分類不能の腎炎	11	43	133	220	189	119	75	46	17	7	860	89	949	20.65	3.47
(%)	(1.3)	(5.0)	(15.5)	(25.6)	(22.0)	(13.8)	(8.7)	(5.3)	(2.0)	(0.8)	(100.0)				
多発性嚢胞腎	24	179	765	1,733	2,069	1,461	730	316	159	101	7,537	661	8,198	21.31	3.16
(%)	(0.3)	(2.4)	(10.1)	(23.0)	(27.5)	(19.4)	(9.7)	(4.2)	(2.1)	(1.3)	(100.0)				
腎硬化症	126	823	2,486	4,183	4,676	3,437	1,940	912	457	377	19,417	2,043	21,460	21.19	3.52
(%)	(0.6)	(4.2)	(12.8)	(21.5)	(24.1)	(17.7)	(10.0)	(4.7)	(2.4)	(1.9)	(100.0)				
悪性高血圧	11	67	179	353	348	298	201	112	73	57	1,699	182	1,881	21.78	3.89
(%)	(0.6)	(3.9)	(10.5)	(20.8)	(20.5)	(17.5)	(11.8)	(6.6)	(4.3)	(3.4)	(100.0)				
糖尿病性腎症	332	1,907	6,904	14,030	17,812	16,098	10,731	6,364	3,473	3,116	80,767	9,740	90,507	22.31	3.80
(%)	(0.4)	(2.4)	(8.5)	(17.4)	(22.1)	(19.9)	(13.3)	(7.9)	(4.3)	(3.9)	(100.0)				
SLE腎炎	26	98	308	371	285	165	93	35	17	20	1,418	132	1,550	19.93	3.44
(%)	(1.8)	(6.9)	(21.7)	(26.2)	(20.1)	(11.6)	(6.6)	(2.5)	(1.2)	(1.4)	(100.0)				
アミロイド腎	8	30	58	74	56	27	20	10	2		285	40	325	19.53	3.24
(%)	(2.8)	(10.5)	(20.4)	(26.0)	(19.6)	(9.5)	(7.0)	(3.5)	(0.7)		(100.0)				
痛風腎	4	22	76	173	196	140	74	36	17	12	750	78	828	21.33	3.39
(%)	(0.5)	(2.9)	(10.1)	(23.1)	(26.1)	(18.7)	(9.9)	(4.8)	(2.3)	(1.6)	(100.0)				
先天性代謝異常による腎不全	3	13	31	30	44	20	11	7	5	5	169	8	177	20.59	3.91
(%)	(1.8)	(7.7)	(18.3)	(17.8)	(26.0)	(11.8)	(6.5)	(4.1)	(3.0)	(3.0)	(100.0)				
腎・尿路結核	1	17	29	29	35	19	13	2			145	13	158	19.87	3.02
(%)	(0.7)	(11.7)	(20.0)	(20.0)	(24.1)	(13.1)	(9.0)	(1.4)			(100.0)				
腎・尿路結石	3	16	46	93	94	66	40	15	10	6	389	41	430	21.13	3.55
(%)	(0.8)	(4.1)	(11.8)	(23.9)	(24.2)	(17.0)	(10.3)	(3.9)	(2.6)	(1.5)	(100.0)				
腎・尿路腫瘍	2	21	82	128	163	121	62	23	10	7	619	67	686	21.08	3.16
(%)	(0.3)	(3.4)	(13.2)	(20.7)	(26.3)	(19.5)	(10.0)	(3.7)	(1.6)	(1.1)	(100.0)				
閉塞性尿路障害	2	30	77	107	106	76	45	19	6	8	476	41	517	20.71	3.48
(%)	(0.4)	(6.3)	(16.2)	(22.5)	(22.3)	(16.0)	(9.5)	(4.0)	(1.3)	(1.7)	(100.0)				
骨髄腫	1	4	27	57	37	32	7	7			172	18	190	20.26	2.68
(%)	(0.6)	(2.3)	(15.7)	(33.1)	(21.5)	(18.6)	(4.1)	(4.1)			(100.0)				
腎形成不全	2	14	51	86	91	64	40	18	12	13	391	28	419	21.33	3.75
(%)	(0.5)	(3.6)	(13.0)	(22.0)	(23.3)	(16.4)	(10.2)	(4.6)	(3.1)	(3.3)	(100.0)				
不明	175	940	2,508	3,937	3,955	2,962	1,587	808	397	322	17,591	2,534	20,125	20.93	3.59
(%)	(1.0)	(5.3)	(14.3)	(22.4)	(22.5)	(16.8)	(9.0)	(4.6)	(2.3)	(1.8)	(100.0)				
再導入	8	75	219	321	312	168	92	50	24	19	1,288	129	1,417	20.52	3.42
(%)	(0.6)	(5.8)	(17.0)	(24.9)	(24.2)	(13.0)	(7.1)	(3.9)	(1.9)	(1.5)	(100.0)				
その他	44	271	716	1,078	1,122	737	437	220	102	81	4,808	706	5,514	20.84	3.57
(%)	(0.9)	(5.6)	(14.9)	(22.4)	(23.3)	(15.3)	(9.1)	(4.6)	(2.1)	(1.7)	(100.0)				
合計	1,311	7,606	24,480	43,156	47,680	36,810	21,913	11,688	5,991	5,164	205,799	23,172	228,971	21.46	3.66
(%)	(0.6)	(3.7)	(11.9)	(21.0)	(23.2)	(17.9)	(10.6)	(5.7)	(2.9)	(2.5)	(100.0)				
記載なし															
(%)															
総計	1,311	7,606	24,480	43,156	47,680	36,810	21,913	11,688	5,991	5,164	205,799	23,172	228,971	21.46	3.66
(%)	(0.6)	(3.7)	(11.9)	(21.0)	(23.2)	(17.9)	(10.6)	(5.7)	(2.9)	(2.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査