

表 2801 Body Mass Index 透析前リン濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前リン濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
3未満	170	727	1,555	1,863	1,617	1,049	535	219	108	81	7,924	1,037	8,961	19.94	3.49
(%)	(2.1)	(9.2)	(19.6)	(23.5)	(20.4)	(13.2)	(6.8)	(2.8)	(1.4)	(1.0)	(100.0)				
3.0～	129	591	1,504	2,386	2,210	1,539	851	400	188	130	9,928	1,052	10,980	20.64	3.49
(%)	(1.3)	(6.0)	(15.1)	(24.0)	(22.3)	(15.5)	(8.6)	(4.0)	(1.9)	(1.3)	(100.0)				
3.5～	147	840	2,476	3,931	4,035	2,947	1,608	781	362	292	17,419	1,599	19,018	20.94	3.51
(%)	(0.8)	(4.8)	(14.2)	(22.6)	(23.2)	(16.9)	(9.2)	(4.5)	(2.1)	(1.7)	(100.0)				
4.0～	166	1,050	3,390	5,823	6,211	4,504	2,501	1,282	602	458	25,987	2,205	28,192	21.14	3.49
(%)	(0.6)	(4.0)	(13.0)	(22.4)	(23.9)	(17.3)	(9.6)	(4.9)	(2.3)	(1.8)	(100.0)				
4.5～	157	1,124	3,662	6,691	7,363	5,535	3,275	1,630	858	637	30,932	2,536	33,468	21.38	3.55
(%)	(0.5)	(3.6)	(11.8)	(21.6)	(23.8)	(17.9)	(10.6)	(5.3)	(2.8)	(2.1)	(100.0)				
5.0～	154	1,028	3,532	6,517	7,484	5,744	3,515	1,782	862	689	31,307	2,502	33,809	21.51	3.55
(%)	(0.5)	(3.3)	(11.3)	(20.8)	(23.9)	(18.3)	(11.2)	(5.7)	(2.8)	(2.2)	(100.0)				
5.5～	139	745	2,908	5,562	6,318	5,020	3,067	1,598	814	731	26,902	2,110	29,012	21.67	3.63
(%)	(0.5)	(2.8)	(10.8)	(20.7)	(23.5)	(18.7)	(11.4)	(5.9)	(3.0)	(2.7)	(100.0)				
6.0～	24	124	496	833	1,097	951	493	310	133	114	4,575	353	4,928	21.76	3.61
(%)	(0.5)	(2.7)	(10.8)	(18.2)	(24.0)	(20.8)	(10.8)	(6.8)	(2.9)	(2.5)	(100.0)				
6.1～	65	414	1,598	2,965	3,418	2,987	1,748	1,023	524	473	15,215	1,290	16,505	21.86	3.69
(%)	(0.4)	(2.7)	(10.5)	(19.5)	(22.5)	(19.6)	(11.5)	(6.7)	(3.4)	(3.1)	(100.0)				
6.5～	59	356	1,264	2,521	3,096	2,468	1,575	943	521	470	13,273	1,104	14,377	22.01	3.76
(%)	(0.4)	(2.7)	(9.5)	(19.0)	(23.3)	(18.6)	(11.9)	(7.1)	(3.9)	(3.5)	(100.0)				
7.0～	22	206	755	1,536	1,817	1,507	996	605	351	316	8,111	749	8,860	22.17	3.82
(%)	(0.3)	(2.5)	(9.3)	(18.9)	(22.4)	(18.6)	(12.3)	(7.5)	(4.3)	(3.9)	(100.0)				
7.5～	23	141	463	870	1,060	892	651	377	215	251	4,943	416	5,359	22.32	4.01
(%)	(0.5)	(2.9)	(9.4)	(17.6)	(21.4)	(18.0)	(13.2)	(7.6)	(4.3)	(5.1)	(100.0)				
8.0～	10	68	241	493	611	549	379	256	136	182	2,925	267	3,192	22.67	4.08
(%)	(0.3)	(2.3)	(8.2)	(16.9)	(20.9)	(18.8)	(13.0)	(8.8)	(4.6)	(6.2)	(100.0)				
8.5～	3	45	158	307	378	350	231	151	79	106	1,808	215	2,023	22.58	4.07
(%)	(0.2)	(2.5)	(8.7)	(17.0)	(20.9)	(19.4)	(12.8)	(8.4)	(4.4)	(5.9)	(100.0)				
9.0～	19	75	235	458	600	483	348	244	189	195	2,846	310	3,156	22.75	4.30
(%)	(0.7)	(2.6)	(8.3)	(16.1)	(21.1)	(17.0)	(12.2)	(8.6)	(6.6)	(6.9)	(100.0)				
合計	1,287	7,534	24,237	42,756	47,315	36,525	21,773	11,601	5,942	5,125	204,095	17,745	221,840	21.47	3.66
(%)	(0.6)	(3.7)	(11.9)	(20.9)	(23.2)	(17.9)	(10.7)	(5.7)	(2.9)	(2.5)	(100.0)				
記載なし	24	72	243	400	365	285	140	87	49	39	1,704	5,427	7,131	21.04	3.72
(%)	(1.4)	(4.2)	(14.3)	(23.5)	(21.4)	(16.7)	(8.2)	(5.1)	(2.9)	(2.3)	(100.0)				
総計	1,311	7,606	24,480	43,156	47,680	36,810	21,913	11,688	5,991	5,164	205,799	23,172	228,971	21.46	3.66
(%)	(0.6)	(3.7)	(11.9)	(21.0)	(23.2)	(17.9)	(10.6)	(5.7)	(2.9)	(2.5)	(100.0)				
平均	4.61	4.76	4.96	5.09	5.19	5.28	5.37	5.50	5.61	5.80	5.20	5.14	5.20		
標準偏差	1.60	1.50	1.41	1.38	1.37	1.37	1.40	1.44	1.53	1.57	1.42	1.54	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査