

表 2802 Body Mass Index PTH値(pg/mL)別(施設血液透析患者 透析回数週3回) PTH測定法:intact-PTH

PTH値	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
20未満	108	525	1,312	2,047	1,968	1,387	749	349	171	130	8,746	766	9,512	20.68	3.52
(%)	(1.2)	(6.0)	(15.0)	(23.4)	(22.5)	(15.9)	(8.6)	(4.0)	(2.0)	(1.5)	(100.0)				
20～	133	644	1,890	2,891	2,816	2,043	1,124	568	306	213	12,628	1,101	13,729	20.88	3.59
(%)	(1.1)	(5.1)	(15.0)	(22.9)	(22.3)	(16.2)	(8.9)	(4.5)	(2.4)	(1.7)	(100.0)				
40～	104	635	1,826	2,950	3,173	2,333	1,252	611	296	232	13,412	1,113	14,525	21.03	3.52
(%)	(0.8)	(4.7)	(13.6)	(22.0)	(23.7)	(17.4)	(9.3)	(4.6)	(2.2)	(1.7)	(100.0)				
60～	116	626	1,851	3,175	3,349	2,426	1,450	709	401	298	14,401	1,113	15,514	21.19	3.61
(%)	(0.8)	(4.3)	(12.9)	(22.0)	(23.3)	(16.8)	(10.1)	(4.9)	(2.8)	(2.1)	(100.0)				
80～	92	560	1,809	3,168	3,510	2,578	1,454	755	385	284	14,595	1,076	15,671	21.28	3.54
(%)	(0.6)	(3.8)	(12.4)	(21.7)	(24.0)	(17.7)	(10.0)	(5.2)	(2.6)	(1.9)	(100.0)				
100～	78	523	1,685	2,974	3,357	2,534	1,490	799	348	350	14,138	1,133	15,271	21.42	3.61
(%)	(0.6)	(3.7)	(11.9)	(21.0)	(23.7)	(17.9)	(10.5)	(5.7)	(2.5)	(2.5)	(100.0)				
120～	63	459	1,574	2,826	3,214	2,446	1,406	815	369	333	13,505	1,081	14,586	21.50	3.60
(%)	(0.5)	(3.4)	(11.7)	(20.9)	(23.8)	(18.1)	(10.4)	(6.0)	(2.7)	(2.5)	(100.0)				
140～	62	374	1,393	2,577	3,025	2,261	1,403	745	379	314	12,533	970	13,503	21.60	3.61
(%)	(0.5)	(3.0)	(11.1)	(20.6)	(24.1)	(18.0)	(11.2)	(5.9)	(3.0)	(2.5)	(100.0)				
160～	52	330	1,258	2,335	2,591	2,124	1,282	694	337	281	11,284	823	12,107	21.64	3.59
(%)	(0.5)	(2.9)	(11.1)	(20.7)	(23.0)	(18.8)	(11.4)	(6.2)	(3.0)	(2.5)	(100.0)				
180～	121	696	2,593	5,085	6,005	4,796	2,971	1,641	793	729	25,430	1,998	27,428	21.78	3.65
(%)	(0.5)	(2.7)	(10.2)	(20.0)	(23.6)	(18.9)	(11.7)	(6.5)	(3.1)	(2.9)	(100.0)				
240～	2	6	27	60	78	42	39	27	13	14	308	30	338	22.19	3.93
(%)	(0.6)	(1.9)	(8.8)	(19.5)	(25.3)	(13.6)	(12.7)	(8.8)	(4.2)	(4.5)	(100.0)				
241～	58	375	1,429	2,822	3,255	2,836	1,737	929	522	458	14,421	1,177	15,598	21.94	3.69
(%)	(0.4)	(2.6)	(9.9)	(19.6)	(22.6)	(19.7)	(12.0)	(6.4)	(3.6)	(3.2)	(100.0)				
300～	35	223	821	1,554	1,888	1,627	968	566	314	275	8,271	721	8,992	21.96	3.74
(%)	(0.4)	(2.7)	(9.9)	(18.8)	(22.8)	(19.7)	(11.7)	(6.8)	(3.8)	(3.3)	(100.0)				
360～	28	132	475	891	1,039	894	573	316	193	170	4,711	413	5,124	22.00	3.83
(%)	(0.6)	(2.8)	(10.1)	(18.9)	(22.1)	(19.0)	(12.2)	(6.7)	(4.1)	(3.6)	(100.0)				
420～	10	104	260	523	610	470	348	220	112	120	2,777	247	3,024	22.09	3.94
(%)	(0.4)	(3.7)	(9.4)	(18.8)	(22.0)	(16.9)	(12.5)	(7.9)	(4.0)	(4.3)	(100.0)				
480～	1	26	70	127	138	111	80	53	29	29	664	72	736	22.06	4.06
(%)	(0.2)	(3.9)	(10.5)	(19.1)	(20.8)	(16.7)	(12.0)	(8.0)	(4.4)	(4.4)	(100.0)				
500～	2	35	128	211	252	179	125	75	41	40	1,088	101	1,189	21.88	3.84
(%)	(0.2)	(3.2)	(11.8)	(19.4)	(23.2)	(16.5)	(11.5)	(6.9)	(3.8)	(3.7)	(100.0)				
540～	5	36	156	224	245	210	125	82	52	55	1,190	120	1,310	21.95	4.07
(%)	(0.4)	(3.0)	(13.1)	(18.8)	(20.6)	(17.6)	(10.5)	(6.9)	(4.4)	(4.6)	(100.0)				
600～	10	59	221	389	434	294	235	126	84	92	1,944	188	2,132	21.97	4.01
(%)	(0.5)	(3.0)	(11.4)	(20.0)	(22.3)	(15.1)	(12.1)	(6.5)	(4.3)	(4.7)	(100.0)				
800～	8	83	173	310	318	224	142	97	51	71	1,477	145	1,622	21.62	4.15
(%)	(0.5)	(5.6)	(11.7)	(21.0)	(21.5)	(15.2)	(9.6)	(6.6)	(3.5)	(4.8)	(100.0)				
合計	1,088	6,451	20,951	37,139	41,265	31,815	18,953	10,177	5,196	4,488	177,523	14,388	191,911	21.48	3.66
(%)	(0.6)	(3.6)	(11.8)	(20.9)	(23.2)	(17.9)	(10.7)	(5.7)	(2.9)	(2.5)	(100.0)				
記載なし	30	172	460	727	722	579	321	163	90	80	3,344	1,331	4,675	21.13	3.72
(%)	(0.9)	(5.1)	(13.8)	(21.7)	(21.6)	(17.3)	(9.6)	(4.9)	(2.7)	(2.4)	(100.0)				
総計	1,118	6,623	21,411	37,866	41,987	32,394	19,274	10,340	5,286	4,568	180,867	15,719	196,586	21.47	3.66
(%)	(0.6)	(3.7)	(11.8)	(20.9)	(23.2)	(17.9)	(10.7)	(5.7)	(2.9)	(2.5)	(100.0)				
平均	139.21	153.45	156.97	163.26	167.60	171.80	178.97	186.82	191.77	210.79	169.62	174.30	169.97		
標準偏差	172.03	181.43	157.06	158.12	157.30	154.68	158.97	164.78	160.67	204.46	160.47	166.91	160.97		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査