

表 2803 Body Mass Index PTH値(pg/mL)別(施設血液透析患者 透析回数週3回) PTH測定法:whole-PTH

PTH値	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
20未満	22	88	207	320	293	235	119	61	30	26	1,401	98	1,499	20.79	3.65
(%)	(1.6)	(6.3)	(14.8)	(22.8)	(20.9)	(16.8)	(8.5)	(4.4)	(2.1)	(1.9)	(100.0)				
20～	20	79	234	373	416	276	198	78	34	34	1,742	106	1,848	21.09	3.58
(%)	(1.1)	(4.5)	(13.4)	(21.4)	(23.9)	(15.8)	(11.4)	(4.5)	(2.0)	(2.0)	(100.0)				
35～	16	57	198	335	321	254	173	80	32	28	1,494	119	1,613	21.21	3.58
(%)	(1.1)	(3.8)	(13.3)	(22.4)	(21.5)	(17.0)	(11.6)	(5.4)	(2.1)	(1.9)	(100.0)				
50～	13	75	242	484	546	441	262	138	67	39	2,307	152	2,459	21.51	3.48
(%)	(0.6)	(3.3)	(10.5)	(21.0)	(23.7)	(19.1)	(11.4)	(6.0)	(2.9)	(1.7)	(100.0)				
75～	6	79	216	391	488	365	218	120	62	55	2,000	148	2,148	21.63	3.64
(%)	(0.3)	(4.0)	(10.8)	(19.6)	(24.4)	(18.3)	(10.9)	(6.0)	(3.1)	(2.8)	(100.0)				
100～	5	52	150	376	384	343	184	84	52	49	1,679	108	1,787	21.70	3.58
(%)	(0.3)	(3.1)	(8.9)	(22.4)	(22.9)	(20.4)	(11.0)	(5.0)	(3.1)	(2.9)	(100.0)				
125～	2	28	111	203	259	235	128	60	34	35	1,095	92	1,187	21.90	3.63
(%)	(0.2)	(2.6)	(10.1)	(18.5)	(23.7)	(21.5)	(11.7)	(5.5)	(3.1)	(3.2)	(100.0)				
150～			4	6	6	9	4	5	1	1	36	1	37	22.37	3.56
(%)			(11.1)	(16.7)	(16.7)	(25.0)	(11.1)	(13.9)	(2.8)	(2.8)	(100.0)				
151～	5	20	88	168	201	156	115	52	32	22	859	75	934	21.84	3.67
(%)	(0.6)	(2.3)	(10.2)	(19.6)	(23.4)	(18.2)	(13.4)	(6.1)	(3.7)	(2.6)	(100.0)				
180～	4	15	64	129	128	124	74	37	23	21	619	47	666	21.89	3.69
(%)	(0.6)	(2.4)	(10.3)	(20.8)	(20.7)	(20.0)	(12.0)	(6.0)	(3.7)	(3.4)	(100.0)				
210～	1	10	47	79	79	88	58	31	18	23	434	38	472	22.31	4.04
(%)	(0.2)	(2.3)	(10.8)	(18.2)	(18.2)	(20.3)	(13.4)	(7.1)	(4.1)	(5.3)	(100.0)				
240～	1	12	44	96	106	76	59	32	13	21	460	66	526	22.03	3.80
(%)	(0.2)	(2.6)	(9.6)	(20.9)	(23.0)	(16.5)	(12.8)	(7.0)	(2.8)	(4.6)	(100.0)				
300～		3	23	24	40	38	13	23	14	8	186	19	205	22.72	4.14
(%)		(1.6)	(12.4)	(12.9)	(21.5)	(20.4)	(7.0)	(12.4)	(7.5)	(4.3)	(100.0)				
350～	1	3	12	23	27	21	10	10	2	3	112	23	135	21.63	3.62
(%)	(0.9)	(2.7)	(10.7)	(20.5)	(24.1)	(18.8)	(8.9)	(8.9)	(1.8)	(2.7)	(100.0)				
400～	1	4	23	27	22	28	16	8	6	4	139	13	152	21.58	3.89
(%)	(0.7)	(2.9)	(16.5)	(19.4)	(15.8)	(20.1)	(11.5)	(5.8)	(4.3)	(2.9)	(100.0)				
500～	1	7	22	34	32	25	21	6	5	2	155	11	166	21.17	3.50
(%)	(0.6)	(4.5)	(14.2)	(21.9)	(20.6)	(16.1)	(13.5)	(3.9)	(3.2)	(1.3)	(100.0)				
合計	98	532	1,685	3,068	3,348	2,714	1,652	825	425	371	14,718	1,116	15,834	21.52	3.65
(%)	(0.7)	(3.6)	(11.4)	(20.8)	(22.7)	(18.4)	(11.2)	(5.6)	(2.9)	(2.5)	(100.0)				
記載なし	5	20	48	72	74	52	23	7	8	5	314	124	438	20.52	3.49
(%)	(1.6)	(6.4)	(15.3)	(22.9)	(23.6)	(16.6)	(7.3)	(2.2)	(2.5)	(1.6)	(100.0)				
総計	103	552	1,733	3,140	3,422	2,766	1,675	832	433	376	15,032	1,240	16,272	21.50	3.64
(%)	(0.7)	(3.7)	(11.5)	(20.9)	(22.8)	(18.4)	(11.1)	(5.5)	(2.9)	(2.5)	(100.0)				
平均	75.09	89.81	103.06	101.64	102.33	106.53	107.71	114.61	120.14	122.41	104.73	118.51	105.70		
標準偏差	111.97	110.87	133.32	116.41	101.89	95.48	100.79	116.87	103.73	98.85	109.20	108.97	109.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査