

表 2808 Body Mass Index 透析前収縮期血圧(mmHg)別（施設血液透析患者 透析回数週3回）

透析前収縮期血圧	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
60未満	2	2	4	9	6	3	3	1		1	31	4	35	20.12	4.08
(%)	(6.5)	(6.5)	(12.9)	(29.0)	(19.4)	(9.7)	(9.7)	(3.2)		(3.2)	(100.0)				
60～	8	38	42	47	56	24	20	9	3	4	251	20	271	19.80	3.89
(%)	(3.2)	(15.1)	(16.7)	(18.7)	(22.3)	(9.6)	(8.0)	(3.6)	(1.2)	(1.6)	(100.0)				
80～	48	189	350	491	454	296	178	71	55	42	2,174	203	2,377	20.43	3.78
(%)	(2.2)	(8.7)	(16.1)	(22.6)	(20.9)	(13.6)	(8.2)	(3.3)	(2.5)	(1.9)	(100.0)				
100～	133	644	1,801	2,645	2,752	1,900	1,118	595	291	212	12,091	1,064	13,155	20.92	3.62
(%)	(1.1)	(5.3)	(14.9)	(21.9)	(22.8)	(15.7)	(9.2)	(4.9)	(2.4)	(1.8)	(100.0)				
120～	291	1,567	4,895	8,441	9,083	6,695	3,815	1,932	962	755	38,436	3,140	41,576	21.21	3.55
(%)	(0.8)	(4.1)	(12.7)	(22.0)	(23.6)	(17.4)	(9.9)	(5.0)	(2.5)	(2.0)	(100.0)				
140～	316	1,998	6,824	12,635	13,936	10,934	6,517	3,313	1,651	1,419	59,543	4,660	64,203	21.49	3.59
(%)	(0.5)	(3.4)	(11.5)	(21.2)	(23.4)	(18.4)	(10.9)	(5.6)	(2.8)	(2.4)	(100.0)				
160～	175	1,347	4,813	8,702	10,172	7,997	4,922	2,707	1,337	1,215	43,387	3,456	46,843	21.69	3.66
(%)	(0.4)	(3.1)	(11.1)	(20.1)	(23.4)	(18.4)	(11.3)	(6.2)	(3.1)	(2.8)	(100.0)				
180～	116	567	1,652	3,091	3,685	3,145	1,903	1,133	671	613	16,576	1,426	18,002	21.94	3.87
(%)	(0.7)	(3.4)	(10.0)	(18.6)	(22.2)	(19.0)	(11.5)	(6.8)	(4.0)	(3.7)	(100.0)				
200～	23	121	456	764	889	751	497	292	192	159	4,144	371	4,515	22.02	3.93
(%)	(0.6)	(2.9)	(11.0)	(18.4)	(21.5)	(18.1)	(12.0)	(7.0)	(4.6)	(3.8)	(100.0)				
220～	2	30	102	170	162	119	91	70	36	42	824	70	894	22.02	4.11
(%)	(0.2)	(3.6)	(12.4)	(20.6)	(19.7)	(14.4)	(11.0)	(8.5)	(4.4)	(5.1)	(100.0)				
合計	1,114	6,503	20,939	36,995	41,195	31,864	19,064	10,123	5,198	4,462	177,457	14,414	191,871	21.48	3.65
(%)	(0.6)	(3.7)	(11.8)	(20.8)	(23.2)	(18.0)	(10.7)	(5.7)	(2.9)	(2.5)	(100.0)				
記載なし	197	1,103	3,541	6,161	6,485	4,946	2,849	1,565	793	702	28,342	8,758	37,100	21.35	3.67
(%)	(0.7)	(3.9)	(12.5)	(21.7)	(22.9)	(17.5)	(10.1)	(5.5)	(2.8)	(2.5)	(100.0)				
総計	1,311	7,606	24,480	43,156	47,680	36,810	21,913	11,688	5,991	5,164	205,799	23,172	228,971	21.46	3.66
(%)	(0.6)	(3.7)	(11.9)	(21.0)	(23.2)	(17.9)	(10.6)	(5.7)	(2.9)	(2.5)	(100.0)				
平均	144.65	147.17	149.28	150.49	151.44	152.66	153.21	154.60	155.65	157.19	151.64	151.61	151.64		
標準偏差	27.80	26.10	24.52	23.73	23.49	23.31	23.54	24.06	24.83	24.82	23.96	24.72	24.02		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査