

表 2837 Body Mass Index PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:intact-PTH

PTH値	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
20未満	9	71	226	384	346	273	139	69	40	20	1,577	106	1,683	20.91	3.42
(%)	(0.6)	(4.5)	(14.3)	(24.4)	(21.9)	(17.3)	(8.8)	(4.4)	(2.5)	(1.3)	(100.0)				
20～	12	84	263	472	489	341	186	102	55	44	2,048	147	2,195	21.18	3.56
(%)	(0.6)	(4.1)	(12.8)	(23.0)	(23.9)	(16.7)	(9.1)	(5.0)	(2.7)	(2.1)	(100.0)				
40～	8	78	284	501	511	406	236	114	60	52	2,250	187	2,437	21.36	3.56
(%)	(0.4)	(3.5)	(12.6)	(22.3)	(22.7)	(18.0)	(10.5)	(5.1)	(2.7)	(2.3)	(100.0)				
60～	11	76	257	521	569	417	285	138	80	61	2,415	177	2,592	21.63	3.63
(%)	(0.5)	(3.1)	(10.6)	(21.6)	(23.6)	(17.3)	(11.8)	(5.7)	(3.3)	(2.5)	(100.0)				
80～	10	81	310	557	603	478	297	164	79	71	2,650	196	2,846	21.59	3.66
(%)	(0.4)	(3.1)	(11.7)	(21.0)	(22.8)	(18.0)	(11.2)	(6.2)	(3.0)	(2.7)	(100.0)				
100～	7	80	288	524	615	465	287	154	96	82	2,598	197	2,795	21.73	3.70
(%)	(0.3)	(3.1)	(11.1)	(20.2)	(23.7)	(17.9)	(11.0)	(5.9)	(3.7)	(3.2)	(100.0)				
120～	9	64	252	500	594	434	265	147	68	67	2,400	181	2,581	21.67	3.58
(%)	(0.4)	(2.7)	(10.5)	(20.8)	(24.8)	(18.1)	(11.0)	(6.1)	(2.8)	(2.8)	(100.0)				
140～	8	53	243	448	474	399	259	139	77	53	2,153	170	2,323	21.73	3.61
(%)	(0.4)	(2.5)	(11.3)	(20.8)	(22.0)	(18.5)	(12.0)	(6.5)	(3.6)	(2.5)	(100.0)				
160～	7	55	217	394	483	355	228	141	60	57	1,997	153	2,150	21.77	3.65
(%)	(0.4)	(2.8)	(10.9)	(19.7)	(24.2)	(17.8)	(11.4)	(7.1)	(3.0)	(2.9)	(100.0)				
180～	16	105	459	945	1,082	855	557	306	172	140	4,637	368	5,005	21.90	3.68
(%)	(0.3)	(2.3)	(9.9)	(20.4)	(23.3)	(18.4)	(12.0)	(6.6)	(3.7)	(3.0)	(100.0)				
240～		2	6	14	19	16	8	3	3	1	72	5	77	21.68	3.26
(%)		(2.8)	(8.3)	(19.4)	(26.4)	(22.2)	(11.1)	(4.2)	(4.2)	(1.4)	(100.0)				
241～	11	61	290	515	618	537	334	196	87	96	2,745	210	2,955	21.98	3.71
(%)	(0.4)	(2.2)	(10.6)	(18.8)	(22.5)	(19.6)	(12.2)	(7.1)	(3.2)	(3.5)	(100.0)				
300～	9	36	153	343	363	271	207	108	58	57	1,605	109	1,714	21.95	3.73
(%)	(0.6)	(2.2)	(9.5)	(21.4)	(22.6)	(16.9)	(12.9)	(6.7)	(3.6)	(3.6)	(100.0)				
360～	4	24	76	215	233	180	116	65	43	26	982	60	1,042	21.89	3.56
(%)	(0.4)	(2.4)	(7.7)	(21.9)	(23.7)	(18.3)	(11.8)	(6.6)	(4.4)	(2.6)	(100.0)				
420～	2	18	47	113	135	102	54	36	17	23	547	44	591	21.89	3.87
(%)	(0.4)	(3.3)	(8.6)	(20.7)	(24.7)	(18.6)	(9.9)	(6.6)	(3.1)	(4.2)	(100.0)				
480～		7	7	33	31	17	18	11	4	7	135	11	146	22.03	4.13
(%)		(5.2)	(5.2)	(24.4)	(23.0)	(12.6)	(13.3)	(8.1)	(3.0)	(5.2)	(100.0)				
500～	2	4	19	43	59	37	22	15	13	8	222	15	237	22.08	3.81
(%)	(0.9)	(1.8)	(8.6)	(19.4)	(26.6)	(16.7)	(9.9)	(6.8)	(5.9)	(3.6)	(100.0)				
540～	1	6	23	55	48	47	33	10	7	7	237	15	252	21.69	3.65
(%)	(0.4)	(2.5)	(9.7)	(23.2)	(20.3)	(19.8)	(13.9)	(4.2)	(3.0)	(3.0)	(100.0)				
600～		6	46	71	86	79	47	22	18	15	390	33	423	22.07	3.75
(%)		(1.5)	(11.8)	(18.2)	(22.1)	(20.3)	(12.1)	(5.6)	(4.6)	(3.8)	(100.0)				
800～	4	15	41	48	81	56	31	16	13	15	320	30	350	21.66	4.11
(%)	(1.3)	(4.7)	(12.8)	(15.0)	(25.3)	(17.5)	(9.7)	(5.0)	(4.1)	(4.7)	(100.0)				
合計	130	926	3,507	6,696	7,439	5,765	3,609	1,956	1,050	902	31,980	2,414	34,394	21.68	3.66
(%)	(0.4)	(2.9)	(11.0)	(20.9)	(23.3)	(18.0)	(11.3)	(6.1)	(3.3)	(2.8)	(100.0)				
記載なし	4	24	70	110	144	99	53	22	12	13	551	183	734	21.23	3.60
(%)	(0.7)	(4.4)	(12.7)	(20.0)	(26.1)	(18.0)	(9.6)	(4.0)	(2.2)	(2.4)	(100.0)				
総計	134	950	3,577	6,806	7,583	5,864	3,662	1,978	1,062	915	32,531	2,597	35,128	21.67	3.66
(%)	(0.4)	(2.9)	(11.0)	(20.9)	(23.3)	(18.0)	(11.3)	(6.1)	(3.3)	(2.8)	(100.0)				
平均	217.76	164.60	167.95	170.55	177.40	178.78	181.42	184.80	192.05	202.53	177.07	179.75	177.26		
標準偏差	414.23	184.60	180.32	159.43	165.20	166.38	153.34	167.20	174.99	184.70	168.20	185.33	169.45		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査