

表 2839 Body Mass Index 透析前ヘモグロビン濃度(g/dL)別 (血液透析濾過患者 透析回数週3回)

透析前ヘモグロビン濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
5未満		1	3	7	4	5	4	2			26		26	21.40	3.28
(%)		(3.8)	(11.5)	(26.9)	(15.4)	(19.2)	(15.4)	(7.7)			(100.0)				
5～	1	1	4	4	3	7	4	2		2	28	3	31	21.94	4.50
(%)	(3.6)	(3.6)	(14.3)	(14.3)	(10.7)	(25.0)	(14.3)	(7.1)		(7.1)	(100.0)				
6～	1	8	14	20	27	16	9	5	3	3	106	8	114	21.03	3.84
(%)	(0.9)	(7.5)	(13.2)	(18.9)	(25.5)	(15.1)	(8.5)	(4.7)	(2.8)	(2.8)	(100.0)				
7～	9	20	58	93	89	64	37	12	9	8	399	46	445	20.70	3.69
(%)	(2.3)	(5.0)	(14.5)	(23.3)	(22.3)	(16.0)	(9.3)	(3.0)	(2.3)	(2.0)	(100.0)				
8～	13	74	207	349	327	253	138	63	37	35	1,496	141	1,637	21.02	3.62
(%)	(0.9)	(4.9)	(13.8)	(23.3)	(21.9)	(16.9)	(9.2)	(4.2)	(2.5)	(2.3)	(100.0)				
9～	28	192	666	1,193	1,280	907	570	287	126	157	5,406	444	5,850	21.39	3.64
(%)	(0.5)	(3.6)	(12.3)	(22.1)	(23.7)	(16.8)	(10.5)	(5.3)	(2.3)	(2.9)	(100.0)				
10～	39	321	1,348	2,546	2,928	2,112	1,306	699	415	329	12,043	904	12,947	21.65	3.63
(%)	(0.3)	(2.7)	(11.2)	(21.1)	(24.3)	(17.5)	(10.8)	(5.8)	(3.4)	(2.7)	(100.0)				
11～	36	275	1,122	2,235	2,463	2,039	1,255	709	363	315	10,812	765	11,577	21.80	3.64
(%)	(0.3)	(2.5)	(10.4)	(20.7)	(22.8)	(18.9)	(11.6)	(6.6)	(3.4)	(2.9)	(100.0)				
12～	14	108	417	822	953	793	490	272	136	106	4,111	271	4,382	21.80	3.60
(%)	(0.3)	(2.6)	(10.1)	(20.0)	(23.2)	(19.3)	(11.9)	(6.6)	(3.3)	(2.6)	(100.0)				
13～	5	28	94	184	216	216	173	84	48	35	1,083	102	1,185	22.35	3.76
(%)	(0.5)	(2.6)	(8.7)	(17.0)	(19.9)	(19.9)	(16.0)	(7.8)	(4.4)	(3.2)	(100.0)				
14～	2	8	27	87	98	99	74	46	30	20	491	32	523	22.82	3.83
(%)	(0.4)	(1.6)	(5.5)	(17.7)	(20.0)	(20.2)	(15.1)	(9.4)	(6.1)	(4.1)	(100.0)				
合計	148	1,036	3,960	7,540	8,388	6,511	4,060	2,181	1,167	1,010	36,001	2,716	38,717	21.67	3.65
(%)	(0.4)	(2.9)	(11.0)	(20.9)	(23.3)	(18.1)	(11.3)	(6.1)	(3.2)	(2.8)	(100.0)				
記載なし	1	10	38	74	74	60	28	16	10	7	318	911	1,229	21.34	3.50
(%)	(0.3)	(3.1)	(11.9)	(23.3)	(23.3)	(18.9)	(8.8)	(5.0)	(3.1)	(2.2)	(100.0)				
総計	149	1,046	3,998	7,614	8,462	6,571	4,088	2,197	1,177	1,017	36,319	3,627	39,946	21.67	3.65
(%)	(0.4)	(2.9)	(11.0)	(21.0)	(23.3)	(18.1)	(11.3)	(6.0)	(3.2)	(2.8)	(100.0)				
平均	10.47	10.62	10.71	10.78	10.82	10.91	10.96	11.02	11.01	10.90	10.84	10.77	10.84		
標準偏差	1.60	1.35	1.25	1.25	1.22	1.27	1.29	1.27	1.26	1.29	1.26	1.29	1.26		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査