

表 2841 Body Mass Index HDL-C濃度(mg/dL)別(血液透析濾過患者 透析回数週3回)

HDL-C濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
20未満		2	8	15	14	12	8	6	5	2	72	9	81	22.11	3.93
(%)		(2.8)	(11.1)	(20.8)	(19.4)	(16.7)	(11.1)	(8.3)	(6.9)	(2.8)	(100.0)				
20～	5	22	87	193	297	337	281	198	145	141	1,706	168	1,874	23.79	4.12
(%)	(0.3)	(1.3)	(5.1)	(11.3)	(17.4)	(19.8)	(16.5)	(11.6)	(8.5)	(8.3)	(100.0)				
30～	14	90	362	781	1,180	1,179	887	565	327	303	5,688	443	6,131	23.03	3.88
(%)	(0.2)	(1.6)	(6.4)	(13.7)	(20.7)	(20.7)	(15.6)	(9.9)	(5.7)	(5.3)	(100.0)				
40～	15	163	608	1,313	1,628	1,409	904	443	227	192	6,902	458	7,360	21.99	3.53
(%)	(0.2)	(2.4)	(8.8)	(19.0)	(23.6)	(20.4)	(13.1)	(6.4)	(3.3)	(2.8)	(100.0)				
50～	21	168	669	1,297	1,444	997	538	284	135	94	5,647	324	5,971	21.27	3.36
(%)	(0.4)	(3.0)	(11.8)	(23.0)	(25.6)	(17.7)	(9.5)	(5.0)	(2.4)	(1.7)	(100.0)				
60～	22	132	541	1,046	886	570	280	110	51	30	3,668	189	3,857	20.56	3.09
(%)	(0.6)	(3.6)	(14.7)	(28.5)	(24.2)	(15.5)	(7.6)	(3.0)	(1.4)	(0.8)	(100.0)				
70～	13	84	341	522	484	243	117	52	24	25	1,905	118	2,023	20.30	3.15
(%)	(0.7)	(4.4)	(17.9)	(27.4)	(25.4)	(12.8)	(6.1)	(2.7)	(1.3)	(1.3)	(100.0)				
80～	8	44	176	288	238	130	58	18	10	5	975	38	1,013	20.11	2.92
(%)	(0.8)	(4.5)	(18.1)	(29.5)	(24.4)	(13.3)	(5.9)	(1.8)	(1.0)	(0.5)	(100.0)				
90～	2	24	87	117	102	42	17	11	4		406	25	431	19.80	2.81
(%)	(0.5)	(5.9)	(21.4)	(28.8)	(25.1)	(10.3)	(4.2)	(2.7)	(1.0)		(100.0)				
100～	2	22	79	105	85	39	20	7	1	3	363	20	383	19.81	2.96
(%)	(0.6)	(6.1)	(21.8)	(28.9)	(23.4)	(10.7)	(5.5)	(1.9)	(0.3)	(0.8)	(100.0)				
合計	102	751	2,958	5,677	6,358	4,958	3,110	1,694	929	795	27,332	1,792	29,124	21.73	3.66
(%)	(0.4)	(2.7)	(10.8)	(20.8)	(23.3)	(18.1)	(11.4)	(6.2)	(3.4)	(2.9)	(100.0)				
記載なし	47	295	1,040	1,937	2,104	1,613	978	503	248	222	8,987	1,835	10,822	21.48	3.59
(%)	(0.5)	(3.3)	(11.6)	(21.6)	(23.4)	(17.9)	(10.9)	(5.6)	(2.8)	(2.5)	(100.0)				
総計	149	1,046	3,998	7,614	8,462	6,571	4,088	2,197	1,177	1,017	36,319	3,627	39,946	21.67	3.65
(%)	(0.4)	(2.9)	(11.0)	(21.0)	(23.3)	(18.1)	(11.3)	(6.0)	(3.2)	(2.8)	(100.0)				
平均	57.63	57.60	57.68	55.42	52.20	48.58	45.87	43.68	41.70	40.36	51.02	48.27	50.85		
標準偏差	19.78	18.81	18.39	17.32	16.79	15.77	14.85	13.94	13.60	13.29	17.13	17.76	17.18		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査