

表 2842 Body Mass Index Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

Non-HDL-C濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
50未満	4	18	78	113	125	81	59	17	11	4	510	19	529	20.94	3.26
(%)	(0.8)	(3.5)	(15.3)	(22.2)	(24.5)	(15.9)	(11.6)	(3.3)	(2.2)	(0.8)	(100.0)				
50～	14	62	279	558	533	402	229	90	60	43	2,270	133	2,403	21.22	3.44
(%)	(0.6)	(2.7)	(12.3)	(24.6)	(23.5)	(17.7)	(10.1)	(4.0)	(2.6)	(1.9)	(100.0)				
70～	13	145	593	1,120	1,178	829	500	256	109	101	4,844	294	5,138	21.31	3.44
(%)	(0.3)	(3.0)	(12.2)	(23.1)	(24.3)	(17.1)	(10.3)	(5.3)	(2.3)	(2.1)	(100.0)				
90～	16	172	650	1,255	1,466	1,060	616	347	176	130	5,888	377	6,265	21.54	3.50
(%)	(0.3)	(2.9)	(11.0)	(21.3)	(24.9)	(18.0)	(10.5)	(5.9)	(3.0)	(2.2)	(100.0)				
110～	23	130	472	987	1,125	889	567	313	185	159	4,850	312	5,162	21.89	3.73
(%)	(0.5)	(2.7)	(9.7)	(20.4)	(23.2)	(18.3)	(11.7)	(6.5)	(3.8)	(3.3)	(100.0)				
130～	8	63	290	536	638	590	377	206	134	112	2,954	195	3,149	22.20	3.78
(%)	(0.3)	(2.1)	(9.8)	(18.1)	(21.6)	(20.0)	(12.8)	(7.0)	(4.5)	(3.8)	(100.0)				
150～	3	47	118	244	323	272	205	135	71	62	1,480	117	1,597	22.45	3.89
(%)	(0.2)	(3.2)	(8.0)	(16.5)	(21.8)	(18.4)	(13.9)	(9.1)	(4.8)	(4.2)	(100.0)				
170～	2	21	55	102	114	121	88	49	36	40	628	54	682	22.64	4.31
(%)	(0.3)	(3.3)	(8.8)	(16.2)	(18.2)	(19.3)	(14.0)	(7.8)	(5.7)	(6.4)	(100.0)				
190～	2	5	27	32	47	37	42	17	19	13	241	21	262	22.69	4.24
(%)	(0.8)	(2.1)	(11.2)	(13.3)	(19.5)	(15.4)	(17.4)	(7.1)	(7.9)	(5.4)	(100.0)				
210～		4	8	20	32	22	13	8	11	7	125	16	141	22.67	4.09
(%)		(3.2)	(6.4)	(16.0)	(25.6)	(17.6)	(10.4)	(6.4)	(8.8)	(5.6)	(100.0)				
合計	85	667	2,570	4,967	5,581	4,303	2,696	1,438	812	671	23,790	1,538	25,328	21.71	3.65
(%)	(0.4)	(2.8)	(10.8)	(20.9)	(23.5)	(18.1)	(11.3)	(6.0)	(3.4)	(2.8)	(100.0)				
記載なし	64	379	1,428	2,647	2,881	2,268	1,392	759	365	346	12,529	2,089	14,618	21.59	3.64
(%)	(0.5)	(3.0)	(11.4)	(21.1)	(23.0)	(18.1)	(11.1)	(6.1)	(2.9)	(2.8)	(100.0)				
総計	149	1,046	3,998	7,614	8,462	6,571	4,088	2,197	1,177	1,017	36,319	3,627	39,946	21.67	3.65
(%)	(0.4)	(2.9)	(11.0)	(21.0)	(23.3)	(18.1)	(11.3)	(6.0)	(3.2)	(2.8)	(100.0)				
平均	103.14	105.80	102.83	103.38	105.33	107.74	110.24	112.95	116.67	118.87	106.88	110.81	107.12		
標準偏差	34.64	33.84	32.83	32.14	32.36	32.87	34.29	33.17	35.93	35.67	33.24	35.18	33.37		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査