

表 2852 Body Mass Index 体重(kg)別（腹膜透析患者 男性のみ）

体重	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
20未満	5	5	4	1	3						18		18	15.71	3.05
(%)	(27.8)	(27.8)	(22.2)	(5.6)	(16.7)						(100.0)				
20～	2	3	3								8	1	9	15.39	1.67
(%)	(25.0)	(37.5)	(37.5)								(100.0)				
30～	4	5	6	2					1		18		18	16.22	3.88
(%)	(22.2)	(27.8)	(33.3)	(11.1)					(5.6)		(100.0)				
40～	1	10	54	95	24	12				1	197	7	204	18.75	1.87
(%)	(0.5)	(5.1)	(27.4)	(48.2)	(12.2)	(6.1)				(0.5)	(100.0)				
50～			18	174	419	228	27	3			869	32	901	21.18	1.56
(%)			(2.1)	(20.0)	(48.2)	(26.2)	(3.1)	(0.3)			(100.0)				
60～				19	257	559	348	92	6	2	1,283	46	1,329	23.41	1.76
(%)				(1.5)	(20.0)	(43.6)	(27.1)	(7.2)	(0.5)	(0.2)	(100.0)				
70～					7	94	273	228	72	16	690	26	716	25.90	1.88
(%)					(1.0)	(13.6)	(39.6)	(33.0)	(10.4)	(2.3)	(100.0)				
80～						1	21	78	96	61	257	16	273	28.62	1.96
(%)						(0.4)	(8.2)	(30.4)	(37.4)	(23.7)	(100.0)				
90～						1	1	4	22	59	87	7	94	30.72	2.05
(%)						(1.1)	(1.1)	(4.6)	(25.3)	(67.8)	(100.0)				
100～									1	10	11	8	19	32.53	1.56
(%)									(9.1)	(90.9)	(100.0)				
110～												10	10		
(%)															
120～												6	6		
(%)															
合計	12	23	85	291	710	895	670	405	198	149	3,438	159	3,597	23.59	3.36
(%)	(0.3)	(0.7)	(2.5)	(8.5)	(20.7)	(26.0)	(19.5)	(11.8)	(5.8)	(4.3)	(100.0)				
記載なし												2,145	2,145		
(%)															
総計	12	23	85	291	710	895	670	405	198	149	3,438	2,304	5,742	23.59	3.36
(%)	(0.3)	(0.7)	(2.5)	(8.5)	(20.7)	(26.0)	(19.5)	(11.8)	(5.8)	(4.3)	(100.0)				
平均	23.09	32.53	43.84	52.40	57.97	63.07	69.26	74.54	81.22	88.49	65.00	74.54	65.42		
標準偏差	13.46	12.71	9.19	5.73	5.69	5.64	5.71	6.52	7.28	8.65	11.88	21.44	12.61		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査