

表 2855 Body Mass Index クレアチニン濃度(mg/dL)別（腹膜透析患者）

クレアチニン濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
2未満		1	2		1	5			1		10	7	17	21.11	3.94
(%)		(10.0)	(20.0)		(10.0)	(50.0)			(10.0)		(100.0)				
2～	3	10	13	18	32	18	13	8	8	3	126	28	154	21.52	4.28
(%)	(2.4)	(7.9)	(10.3)	(14.3)	(25.4)	(14.3)	(10.3)	(6.3)	(6.3)	(2.4)	(100.0)				
4～	8	16	41	68	123	103	57	40	28	23	507	68	575	22.34	4.07
(%)	(1.6)	(3.2)	(8.1)	(13.4)	(24.3)	(20.3)	(11.2)	(7.9)	(5.5)	(4.5)	(100.0)				
6～	9	18	55	139	232	216	149	91	46	24	979	154	1,133	22.56	3.53
(%)	(0.9)	(1.8)	(5.6)	(14.2)	(23.7)	(22.1)	(15.2)	(9.3)	(4.7)	(2.5)	(100.0)				
8～	1	13	53	156	274	297	179	116	49	33	1,171	178	1,349	22.82	3.35
(%)	(0.1)	(1.1)	(4.5)	(13.3)	(23.4)	(25.4)	(15.3)	(9.9)	(4.2)	(2.8)	(100.0)				
10～	4	6	28	121	197	243	194	106	50	44	993	122	1,115	23.36	3.45
(%)	(0.4)	(0.6)	(2.8)	(12.2)	(19.8)	(24.5)	(19.5)	(10.7)	(5.0)	(4.4)	(100.0)				
12～	2	2	18	57	158	189	149	88	56	33	752	96	848	23.77	3.37
(%)	(0.3)	(0.3)	(2.4)	(7.6)	(21.0)	(25.1)	(19.8)	(11.7)	(7.4)	(4.4)	(100.0)				
14～		2	6	28	72	95	87	63	38	31	422	66	488	24.46	3.43
(%)		(0.5)	(1.4)	(6.6)	(17.1)	(22.5)	(20.6)	(14.9)	(9.0)	(7.3)	(100.0)				
16～		1	3	22	32	62	54	31	19	23	247	48	295	24.44	3.65
(%)		(0.4)	(1.2)	(8.9)	(13.0)	(25.1)	(21.9)	(12.6)	(7.7)	(9.3)	(100.0)				
合計	27	69	219	609	1,121	1,228	882	543	295	214	5,207	767	5,974	23.14	3.60
(%)	(0.5)	(1.3)	(4.2)	(11.7)	(21.5)	(23.6)	(16.9)	(10.4)	(5.7)	(4.1)	(100.0)				
記載なし		2	9	18	19	33	26	18	7	6	138	2,829	2,967	23.30	3.64
(%)		(1.4)	(6.5)	(13.0)	(13.8)	(23.9)	(18.8)	(13.0)	(5.1)	(4.3)	(100.0)				
総計	27	71	228	627	1,140	1,261	908	561	302	220	5,345	3,596	8,941	23.15	3.60
(%)	(0.5)	(1.3)	(4.3)	(11.7)	(21.3)	(23.6)	(17.0)	(10.5)	(5.7)	(4.1)	(100.0)				
平均	7.01	7.14	8.11	9.26	9.49	10.07	10.50	10.54	10.58	11.07	9.91	9.76	9.89		
標準偏差	2.71	3.30	3.08	3.18	3.29	3.49	3.41	3.58	3.86	3.90	3.50	3.68	3.53		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査