

表 2857 Body Mass Index CRP濃度(mg/dL)別（腹膜透析患者）

CRP濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
0.2未満	15	40	115	366	628	669	451	259	124	72	2,739	328	3,067	22.79	3.40
(%)	(0.5)	(1.5)	(4.2)	(13.4)	(22.9)	(24.4)	(16.5)	(9.5)	(4.5)	(2.6)	(100.0)				
0.2～	3	7	32	76	164	190	163	113	69	53	870	125	995	23.85	3.71
(%)	(0.3)	(0.8)	(3.7)	(8.7)	(18.9)	(21.8)	(18.7)	(13.0)	(7.9)	(6.1)	(100.0)				
0.5～	2	7	12	45	55	90	62	51	31	24	379	70	449	23.74	3.86
(%)	(0.5)	(1.8)	(3.2)	(11.9)	(14.5)	(23.7)	(16.4)	(13.5)	(8.2)	(6.3)	(100.0)				
1.0～		3	10	29	52	64	48	20	17	18	261	47	308	23.46	3.72
(%)		(1.1)	(3.8)	(11.1)	(19.9)	(24.5)	(18.4)	(7.7)	(6.5)	(6.9)	(100.0)				
2.0～	1	4	8	15	42	37	19	21	10	9	166	36	202	23.14	3.96
(%)	(0.6)	(2.4)	(4.8)	(9.0)	(25.3)	(22.3)	(11.4)	(12.7)	(6.0)	(5.4)	(100.0)				
4.0～		1	3	7	17	16	11	6	1	8	70	13	83	23.40	4.02
(%)		(1.4)	(4.3)	(10.0)	(24.3)	(22.9)	(15.7)	(8.6)	(1.4)	(11.4)	(100.0)				
6.0～		1	3	3	8	7	3	2	2	1	30	10	40	22.19	3.75
(%)		(3.3)	(10.0)	(10.0)	(26.7)	(23.3)	(10.0)	(6.7)	(6.7)	(3.3)	(100.0)				
8.0～				4	6	2	2	2	1	1	18	6	24	22.74	3.66
(%)				(22.2)	(33.3)	(11.1)	(11.1)	(11.1)	(5.6)	(5.6)	(100.0)				
10.0～			3	4	6	2	11	3			29	5	34	22.63	3.21
(%)			(10.3)	(13.8)	(20.7)	(6.9)	(37.9)	(10.3)			(100.0)				
15.0～		1			1		2		1		5	2	7	22.78	5.22
(%)		(20.0)			(20.0)		(40.0)		(20.0)		(100.0)				
合計	21	64	186	549	979	1,077	772	477	256	186	4,567	642	5,209	23.13	3.58
(%)	(0.5)	(1.4)	(4.1)	(12.0)	(21.4)	(23.6)	(16.9)	(10.4)	(5.6)	(4.1)	(100.0)				
記載なし	6	7	42	78	161	184	136	84	46	34	778	2,954	3,732	23.25	3.69
(%)	(0.8)	(0.9)	(5.4)	(10.0)	(20.7)	(23.7)	(17.5)	(10.8)	(5.9)	(4.4)	(100.0)				
総計	27	71	228	627	1,140	1,261	908	561	302	220	5,345	3,596	8,941	23.15	3.60
(%)	(0.5)	(1.3)	(4.3)	(11.7)	(21.3)	(23.6)	(17.0)	(10.5)	(5.7)	(4.1)	(100.0)				
平均	0.27	0.88	0.70	0.55	0.62	0.48	0.68	0.58	0.60	0.77	0.60	0.87	0.63		
標準偏差	0.53	2.68	1.76	1.53	1.63	1.11	2.26	1.47	1.86	1.29	1.65	2.09	1.71		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査