

表 2860 Body Mass Index PTH値(pg/mL)別（腹膜透析患者）PTH測定法:intact-PTH

PTH値	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
20未満	1	2	6	8	33	22	19	4	9	4	108	15	123	22.76	3.87
(%)	(0.9)	(1.9)	(5.6)	(7.4)	(30.6)	(20.4)	(17.6)	(3.7)	(8.3)	(3.7)	(100.0)				
20～	4	6	12	33	47	66	44	22	8	8	250	31	281	22.64	3.64
(%)	(1.6)	(2.4)	(4.8)	(13.2)	(18.8)	(26.4)	(17.6)	(8.8)	(3.2)	(3.2)	(100.0)				
40～		4	10	28	45	55	49	14	14	11	230	38	268	23.13	3.61
(%)		(1.7)	(4.3)	(12.2)	(19.6)	(23.9)	(21.3)	(6.1)	(6.1)	(4.8)	(100.0)				
60～		3	16	34	63	65	44	23	13	13	274	41	315	22.99	3.61
(%)		(1.1)	(5.8)	(12.4)	(23.0)	(23.7)	(16.1)	(8.4)	(4.7)	(4.7)	(100.0)				
80～	2	5	12	33	52	68	45	36	15	7	275	34	309	23.01	3.52
(%)	(0.7)	(1.8)	(4.4)	(12.0)	(18.9)	(24.7)	(16.4)	(13.1)	(5.5)	(2.5)	(100.0)				
100～	1	3	10	33	58	79	45	26	9	11	275	41	316	23.01	3.36
(%)	(0.4)	(1.1)	(3.6)	(12.0)	(21.1)	(28.7)	(16.4)	(9.5)	(3.3)	(4.0)	(100.0)				
120～	1	3	4	32	47	77	45	21	17	12	259	46	305	23.37	3.54
(%)	(0.4)	(1.2)	(1.5)	(12.4)	(18.1)	(29.7)	(17.4)	(8.1)	(6.6)	(4.6)	(100.0)				
140～		3	8	29	50	62	43	25	15	11	246	39	285	23.45	3.54
(%)		(1.2)	(3.3)	(11.8)	(20.3)	(25.2)	(17.5)	(10.2)	(6.1)	(4.5)	(100.0)				
160～	1	1	15	27	51	50	34	29	17	8	233	43	276	23.16	3.58
(%)	(0.4)	(0.4)	(6.4)	(11.6)	(21.9)	(21.5)	(14.6)	(12.4)	(7.3)	(3.4)	(100.0)				
180～	3	6	22	60	127	151	87	61	38	25	580	74	654	23.28	3.54
(%)	(0.5)	(1.0)	(3.8)	(10.3)	(21.9)	(26.0)	(15.0)	(10.5)	(6.6)	(4.3)	(100.0)				
240～				2	2	2		2			8	2	10	22.57	3.51
(%)				(25.0)	(25.0)	(25.0)		(25.0)			(100.0)				
241～	3	4	14	37	82	84	68	52	25	13	382	49	431	23.45	3.60
(%)	(0.8)	(1.0)	(3.7)	(9.7)	(21.5)	(22.0)	(17.8)	(13.6)	(6.5)	(3.4)	(100.0)				
300～		4	10	33	77	71	37	27	24	12	295	36	331	23.19	3.56
(%)		(1.4)	(3.4)	(11.2)	(26.1)	(24.1)	(12.5)	(9.2)	(8.1)	(4.1)	(100.0)				
360～	1	2	10	22	48	24	30	17	9	10	173	27	200	22.96	3.89
(%)	(0.6)	(1.2)	(5.8)	(12.7)	(27.7)	(13.9)	(17.3)	(9.8)	(5.2)	(5.8)	(100.0)				
420～		4	7	19	17	32	15	12	6	6	118	21	139	22.80	3.75
(%)		(3.4)	(5.9)	(16.1)	(14.4)	(27.1)	(12.7)	(10.2)	(5.1)	(5.1)	(100.0)				
480～			3	4	5	9	5	5	2	4	37	6	43	23.70	4.15
(%)			(8.1)	(10.8)	(13.5)	(24.3)	(13.5)	(13.5)	(5.4)	(10.8)	(100.0)				
500～		1	6	7	9	9	7	8	4	4	46	8	54	24.41	3.89
(%)		(2.2)	(13.0)	(15.2)	(19.6)	(15.2)	(17.4)	(8.7)	(8.7)	(100.0)					
540～		3	5	23	9	6	4	4	2	4	56	17	73	22.89	3.66
(%)		(5.4)	(8.9)	(41.1)	(16.1)	(10.7)	(7.1)	(3.6)	(7.1)	(100.0)					
600～	1	2	3	19	21	15	9	8	6	7	91	14	105	22.82	4.29
(%)	(1.1)	(2.2)	(3.3)	(20.9)	(23.1)	(16.5)	(9.9)	(8.8)	(6.6)	(7.7)	(100.0)				
800～	3	4	6	14	22	14	16	8	3	4	94	16	110	22.14	4.36
(%)	(3.2)	(4.3)	(6.4)	(14.9)	(23.4)	(14.9)	(17.0)	(8.5)	(3.2)	(4.3)	(100.0)				
合計	21	56	172	478	877	964	648	404	236	174	4,030	598	4,628	23.13	3.64
(%)	(0.5)	(1.4)	(4.3)	(11.9)	(21.8)	(23.9)	(16.1)	(10.0)	(5.9)	(4.3)	(100.0)				
記載なし	1	3	11	27	39	53	36	28	12	6	216	160	376	23.08	3.53
(%)	(0.5)	(1.4)	(5.1)	(12.5)	(18.1)	(24.5)	(16.7)	(13.0)	(5.6)	(2.8)	(100.0)				
総計	22	59	183	505	916	1,017	684	432	248	180	4,246	758	5,004	23.12	3.63
(%)	(0.5)	(1.4)	(4.3)	(11.9)	(21.6)	(24.0)	(16.1)	(10.2)	(5.8)	(4.2)	(100.0)				
平均	291.33	267.91	231.46	239.85	233.52	201.26	213.01	225.51	220.30	251.53	223.15	228.79	223.88		
標準偏差	315.91	324.31	241.50	257.03	259.47	184.46	253.04	183.41	173.08	263.38	231.63	213.15	229.31		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査