

表 2866 Body Mass Index 収縮期血圧(mmHg)別（腹膜透析患者）

収縮期血圧	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
60未満	1										1		1	12.90	
(%)	(100.0)										(100.0)				
60～	2		2	2	3	3			1		13		13	20.05	4.13
(%)	(15.4)		(15.4)	(15.4)	(23.1)	(23.1)			(7.7)		(100.0)				
80～	6	8	14	13	29	18	16	8	1	1	114	10	124	20.80	4.09
(%)	(5.3)	(7.0)	(12.3)	(11.4)	(25.4)	(15.8)	(14.0)	(7.0)	(0.9)	(0.9)	(100.0)				
100～	12	24	40	94	123	144	79	39	26	19	600	41	641	22.09	3.84
(%)	(2.0)	(4.0)	(6.7)	(15.7)	(20.5)	(24.0)	(13.2)	(6.5)	(4.3)	(3.2)	(100.0)				
120～	3	24	65	167	326	337	252	139	74	50	1,437	79	1,516	22.99	3.50
(%)	(0.2)	(1.7)	(4.5)	(11.6)	(22.7)	(23.5)	(17.5)	(9.7)	(5.1)	(3.5)	(100.0)				
140～	1	6	48	162	315	341	271	173	87	70	1,474	105	1,579	23.51	3.45
(%)	(0.1)	(0.4)	(3.3)	(11.0)	(21.4)	(23.1)	(18.4)	(11.7)	(5.9)	(4.7)	(100.0)				
160～		1	16	67	128	153	127	89	54	33	668	53	721	23.93	3.46
(%)		(0.1)	(2.4)	(10.0)	(19.2)	(22.9)	(19.0)	(13.3)	(8.1)	(4.9)	(100.0)				
180～		1	1	20	38	54	31	20	13	16	194	23	217	24.03	3.56
(%)		(0.5)	(0.5)	(10.3)	(19.6)	(27.8)	(16.0)	(10.3)	(6.7)	(8.2)	(100.0)				
200～				6	4	6	5	4	1	2	28	3	31	23.63	3.41
(%)				(21.4)	(14.3)	(21.4)	(17.9)	(14.3)	(3.6)	(7.1)	(100.0)				
220～					1	1	4	1	3		10		10	25.49	2.38
(%)					(10.0)	(10.0)	(40.0)	(10.0)	(30.0)		(100.0)				
合計	25	64	186	531	967	1,057	785	473	260	191	4,539	314	4,853	23.17	3.61
(%)	(0.6)	(1.4)	(4.1)	(11.7)	(21.3)	(23.3)	(17.3)	(10.4)	(5.7)	(4.2)	(100.0)				
記載なし	2	7	42	96	173	204	123	88	42	29	806	3,282	4,088	23.03	3.51
(%)	(0.2)	(0.9)	(5.2)	(11.9)	(21.5)	(25.3)	(15.3)	(10.9)	(5.2)	(3.6)	(100.0)				
総計	27	71	228	627	1,140	1,261	908	561	302	220	5,345	3,596	8,941	23.15	3.60
(%)	(0.5)	(1.3)	(4.3)	(11.7)	(21.3)	(23.6)	(17.0)	(10.5)	(5.7)	(4.1)	(100.0)				
平均	104.04	119.09	129.88	138.23	139.68	141.39	143.06	145.04	145.51	146.29	140.78	144.40	141.01		
標準偏差	19.42	20.48	22.27	22.98	21.91	22.42	22.38	21.64	22.78	22.09	22.79	23.87	22.88		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査