

表 2883 透析前BUN(mg/dL) 透析前リン濃度(mg/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
3未満	225	815	1,483	1,700	1,243	804	377	149	57	17	7	6	6,883	6	6,889	46.35	16.68
(%)	(3.3)	(11.8)	(21.5)	(24.7)	(18.1)	(11.7)	(5.5)	(2.2)	(0.8)	(0.2)	(0.1)	(0.1)	(100.0)				
3.0～	60	460	1,491	2,129	1,922	1,245	600	213	75	22	5	1	8,223	13	8,236	50.56	15.14
(%)	(0.7)	(5.6)	(18.1)	(25.9)	(23.4)	(15.1)	(7.3)	(2.6)	(0.9)	(0.3)	(0.1)	(0.0)	(100.0)				
3.5～	38	400	1,865	3,597	3,789	2,627	1,222	463	130	23	6	7	14,167	27	14,194	53.49	14.49
(%)	(0.3)	(2.8)	(13.2)	(25.4)	(26.7)	(18.5)	(8.6)	(3.3)	(0.9)	(0.2)	(0.0)	(0.0)	(100.0)				
4.0～	41	329	1,878	4,774	6,154	4,697	2,425	827	232	46	16	4	21,423	29	21,452	56.19	13.83
(%)	(0.2)	(1.5)	(8.8)	(22.3)	(28.7)	(21.9)	(11.3)	(3.9)	(1.1)	(0.2)	(0.1)	(0.0)	(100.0)				
4.5～	25	182	1,455	4,715	7,587	6,557	3,714	1,297	380	89	27	5	26,033	45	26,078	59.01	13.58
(%)	(0.1)	(0.7)	(5.6)	(18.1)	(29.1)	(25.2)	(14.3)	(5.0)	(1.5)	(0.3)	(0.1)	(0.0)	(100.0)				
5.0～	26	119	938	3,941	7,381	7,263	4,421	1,729	587	141	27	18	26,591	33	26,624	61.41	13.70
(%)	(0.1)	(0.4)	(3.5)	(14.8)	(27.8)	(27.3)	(16.6)	(6.5)	(2.2)	(0.5)	(0.1)	(0.1)	(100.0)				
5.5～	14	62	559	2,703	5,785	6,730	4,430	1,986	628	132	40	18	23,087	35	23,122	63.62	13.65
(%)	(0.1)	(0.3)	(2.4)	(11.7)	(25.1)	(29.2)	(19.2)	(8.6)	(2.7)	(0.6)	(0.2)	(0.1)	(100.0)				
6.0～		8	105	376	926	1,139	831	398	128	35	7	6	3,959	7	3,966	65.02	13.91
(%)		(0.2)	(2.7)	(9.5)	(23.4)	(28.8)	(21.0)	(10.1)	(3.2)	(0.9)	(0.2)	(0.2)	(100.0)				
6.1～	6	23	199	1,134	3,003	3,818	2,892	1,406	516	107	31	13	13,148	17	13,165	65.98	13.68
(%)	(0.0)	(0.2)	(1.5)	(8.6)	(22.8)	(29.0)	(22.0)	(10.7)	(3.9)	(0.8)	(0.2)	(0.1)	(100.0)				
6.5～	8	18	151	758	2,187	3,276	2,675	1,536	538	143	40	15	11,345	17	11,362	67.98	14.04
(%)	(0.1)	(0.2)	(1.3)	(6.7)	(19.3)	(28.9)	(23.6)	(13.5)	(4.7)	(1.3)	(0.4)	(0.1)	(100.0)				
7.0～	6	8	62	416	1,167	1,910	1,822	1,074	439	124	34	10	7,072	8	7,080	69.83	14.31
(%)	(0.1)	(0.1)	(0.9)	(5.9)	(16.5)	(27.0)	(25.8)	(15.2)	(6.2)	(1.8)	(0.5)	(0.1)	(100.0)				
7.5～	2	5	27	153	605	1,086	1,117	685	338	129	25	20	4,192	6	4,198	72.22	14.77
(%)	(0.0)	(0.1)	(0.6)	(3.6)	(14.4)	(25.9)	(26.6)	(16.3)	(8.1)	(3.1)	(0.6)	(0.5)	(100.0)				
8.0～	3	3	19	88	309	552	638	503	235	90	26	16	2,482	3	2,485	74.11	15.57
(%)	(0.1)	(0.1)	(0.8)	(3.5)	(12.4)	(22.2)	(25.7)	(20.3)	(9.5)	(3.6)	(1.0)	(0.6)	(100.0)				
8.5～		2	13	75	188	325	406	312	172	68	28	12	1,601		1,601	74.72	16.24
(%)		(0.1)	(0.8)	(4.7)	(11.7)	(20.3)	(25.4)	(19.5)	(10.7)	(4.2)	(1.7)	(0.7)	(100.0)				
9.0～	3	7	27	89	201	423	608	558	321	172	73	52	2,534	3	2,537	78.64	18.48
(%)	(0.1)	(0.3)	(1.1)	(3.5)	(7.9)	(16.7)	(24.0)	(22.0)	(12.7)	(6.8)	(2.9)	(2.1)	(100.0)				
合計	457	2,441	10,272	26,648	42,447	42,452	28,178	13,136	4,776	1,338	392	203	172,740	249	172,989	60.96	15.62
(%)	(0.3)	(1.4)	(5.9)	(15.4)	(24.6)	(24.6)	(16.3)	(7.6)	(2.8)	(0.8)	(0.2)	(0.1)	(100.0)				
記載なし	22	39	86	163	212	191	144	76	27	10	2	1	973	4,247	5,220	58.03	18.47
(%)	(2.3)	(4.0)	(8.8)	(16.8)	(21.8)	(19.6)	(14.8)	(7.8)	(2.8)	(1.0)	(0.2)	(0.1)	(100.0)				
総計	479	2,480	10,358	26,811	42,659	42,643	28,322	13,212	4,803	1,348	394	204	173,713	4,496	178,209	60.94	15.64
(%)	(0.3)	(1.4)	(6.0)	(15.4)	(24.6)	(24.5)	(16.3)	(7.6)	(2.8)	(0.8)	(0.2)	(0.1)	(100.0)				
平均	3.26	3.49	4.09	4.62	5.05	5.39	5.70	6.05	6.34	6.84	7.05	7.55	5.22	5.13	5.22		
標準偏差	1.65	1.21	1.17	1.17	1.19	1.26	1.38	1.51	1.68	2.01	2.12	2.60	1.43	1.29	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査