

表 2885 透析前BUN(mg/dL) PTH値(pg/mL)別(施設血液透析患者 透析回数週3回 透析歴2年以上) PTH測定法:whole-PTH

PTH値	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
20未満	5	28	113	224	290	281	149	67	24	13	1	3	1,198	1	1,199	57.90	16.25
(%)	(0.4)	(2.3)	(9.4)	(18.7)	(24.2)	(23.5)	(12.4)	(5.6)	(2.0)	(1.1)	(0.1)	(0.3)	(100.0)				
20～	13	31	132	250	404	298	237	84	31	7	4	1	1,492		1,492	58.13	15.94
(%)	(0.9)	(2.1)	(8.8)	(16.8)	(27.1)	(20.0)	(15.9)	(5.6)	(2.1)	(0.5)	(0.3)	(0.1)	(100.0)				
35～	4	28	112	245	361	281	181	83	26	6	1	1	1,329	1	1,330	58.08	15.30
(%)	(0.3)	(2.1)	(8.4)	(18.4)	(27.2)	(21.1)	(13.6)	(6.2)	(2.0)	(0.5)	(0.1)	(0.1)	(100.0)				
50～	6	32	143	348	520	452	289	123	54	8	1		1,976	1	1,977	59.15	15.11
(%)	(0.3)	(1.6)	(7.2)	(17.6)	(26.3)	(22.9)	(14.6)	(6.2)	(2.7)	(0.4)	(0.1)		(100.0)				
75～	4	21	85	296	479	398	295	104	39	9	2		1,732	4	1,736	59.98	14.51
(%)	(0.2)	(1.2)	(4.9)	(17.1)	(27.7)	(23.0)	(17.0)	(6.0)	(2.3)	(0.5)	(0.1)		(100.0)				
100～	3	19	60	239	359	353	231	101	34	8	1	1	1,409		1,409	60.77	14.78
(%)	(0.2)	(1.3)	(4.3)	(17.0)	(25.5)	(25.1)	(16.4)	(7.2)	(2.4)	(0.6)	(0.1)	(0.1)	(100.0)				
125～	1	7	49	123	248	229	158	61	34	3	1	1	915		915	61.56	14.81
(%)	(0.1)	(0.8)	(5.4)	(13.4)	(27.1)	(25.0)	(17.3)	(6.7)	(3.7)	(0.3)	(0.1)	(0.1)	(100.0)				
150～			3	7	6	4	6	1	1				28		28	57.79	15.19
(%)			(10.7)	(25.0)	(21.4)	(14.3)	(21.4)	(3.6)	(3.6)				(100.0)				
151～		7	32	108	191	180	143	47	25	9	2	1	745	2	747	62.25	15.04
(%)		(0.9)	(4.3)	(14.5)	(25.6)	(24.2)	(19.2)	(6.3)	(3.4)	(1.2)	(0.3)	(0.1)	(100.0)				
180～	1	6	21	89	120	118	111	36	17	6	2	1	528		528	62.25	15.95
(%)	(0.2)	(1.1)	(4.0)	(16.9)	(22.7)	(22.3)	(21.0)	(6.8)	(3.2)	(1.1)	(0.4)	(0.2)	(100.0)				
210～	1	3	16	52	87	104	57	30	11	3	2		366		366	62.14	15.22
(%)	(0.3)	(0.8)	(4.4)	(14.2)	(23.8)	(28.4)	(15.6)	(8.2)	(3.0)	(0.8)	(0.5)		(100.0)				
240～		2	18	48	101	106	86	40	10	7	1		419		419	63.71	15.03
(%)		(0.5)	(4.3)	(11.5)	(24.1)	(25.3)	(20.5)	(9.5)	(2.4)	(1.7)	(0.2)		(100.0)				
300～			3	23	29	61	35	13	7	6			177		177	65.30	14.71
(%)			(1.7)	(13.0)	(16.4)	(34.5)	(19.8)	(7.3)	(4.0)	(3.4)			(100.0)				
350～		1	5	17	17	33	21	10	1				105		105	62.26	14.01
(%)		(1.0)	(4.8)	(16.2)	(16.2)	(31.4)	(20.0)	(9.5)	(1.0)				(100.0)				
400～			5	19	31	20	22	17	7	2	1		124		124	64.75	17.07
(%)			(4.0)	(15.3)	(25.0)	(16.1)	(17.7)	(13.7)	(5.6)	(1.6)	(0.8)		(100.0)				
500～			9	9	34	38	23	21	8	8	1	1	152		152	68.11	18.04
(%)			(5.9)	(5.9)	(22.4)	(25.0)	(15.1)	(13.8)	(5.3)	(5.3)	(0.7)	(0.7)	(100.0)				
合計	38	185	806	2,097	3,277	2,956	2,044	838	329	95	20	10	12,695	9	12,704	60.09	15.40
(%)	(0.3)	(1.5)	(6.3)	(16.5)	(25.8)	(23.3)	(16.1)	(6.6)	(2.6)	(0.7)	(0.2)	(0.1)	(100.0)				
記載なし	1	12	29	54	42	33	22	2	6				201	111	312	51.96	15.92
(%)	(0.5)	(6.0)	(14.4)	(26.9)	(20.9)	(16.4)	(10.9)	(1.0)	(3.0)				(100.0)				
総計	39	197	835	2,151	3,319	2,989	2,066	840	335	95	20	10	12,896	120	13,016	59.96	15.44
(%)	(0.3)	(1.5)	(6.5)	(16.7)	(25.7)	(23.2)	(16.0)	(6.5)	(2.6)	(0.7)	(0.2)	(0.1)	(100.0)				
平均	56.18	70.57	85.32	94.92	101.17	111.36	113.57	123.22	129.17	215.54	154.40	145.30	106.08	87.89	106.06		
標準偏差	49.36	58.72	102.96	87.96	102.16	114.77	103.55	120.54	157.18	344.80	143.57	220.54	110.71	55.00	110.69		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査