

表 2886 透析前BUN(mg/dL) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前ヘモグロビン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
5未満	1	4	10	28	45	57	34	17	7	4	2		209	1	210	62.54	16.93
(%)	(0.5)	(1.9)	(4.8)	(13.4)	(21.5)	(27.3)	(16.3)	(8.1)	(3.3)	(1.9)	(1.0)		(100.0)				
5～		8	12	46	61	59	29	20	10	4	1	1	251		251	60.63	17.26
(%)		(3.2)	(4.8)	(18.3)	(24.3)	(23.5)	(11.6)	(8.0)	(4.0)	(1.6)	(0.4)	(0.4)	(100.0)				
6～	8	36	76	119	135	106	68	33	21	7	4	2	615	2	617	56.17	19.21
(%)	(1.3)	(5.9)	(12.4)	(19.3)	(22.0)	(17.2)	(11.1)	(5.4)	(3.4)	(1.1)	(0.7)	(0.3)	(100.0)				
7～	31	124	292	448	556	475	278	140	56	21	6	10	2,437	8	2,445	56.02	18.15
(%)	(1.3)	(5.1)	(12.0)	(18.4)	(22.8)	(19.5)	(11.4)	(5.7)	(2.3)	(0.9)	(0.2)	(0.4)	(100.0)				
8～	58	296	904	1,701	1,970	1,639	1,096	530	193	56	22	16	8,481	16	8,497	57.00	17.17
(%)	(0.7)	(3.5)	(10.7)	(20.1)	(23.2)	(19.3)	(12.9)	(6.2)	(2.3)	(0.7)	(0.3)	(0.2)	(100.0)				
9～	117	566	2,253	4,940	7,195	6,485	4,087	1,865	762	192	63	40	28,565	43	28,608	59.16	16.05
(%)	(0.4)	(2.0)	(7.9)	(17.3)	(25.2)	(22.7)	(14.3)	(6.5)	(2.7)	(0.7)	(0.2)	(0.1)	(100.0)				
10～	134	714	3,328	9,266	15,055	14,869	9,721	4,456	1,545	434	117	57	59,696	73	59,769	60.94	15.25
(%)	(0.2)	(1.2)	(5.6)	(15.5)	(25.2)	(24.9)	(16.3)	(7.5)	(2.6)	(0.7)	(0.2)	(0.1)	(100.0)				
11～	80	479	2,355	6,862	11,806	12,480	8,425	3,958	1,345	388	120	53	48,351	64	48,415	62.03	15.18
(%)	(0.2)	(1.0)	(4.9)	(14.2)	(24.4)	(25.8)	(17.4)	(8.2)	(2.8)	(0.8)	(0.2)	(0.1)	(100.0)				
12～	31	164	776	2,360	4,023	4,429	3,161	1,483	559	158	37	10	17,191	17	17,208	62.62	15.23
(%)	(0.2)	(1.0)	(4.5)	(13.7)	(23.4)	(25.8)	(18.4)	(8.6)	(3.3)	(0.9)	(0.2)	(0.1)	(100.0)				
13～	10	47	165	535	966	1,100	821	379	171	40	11	8	4,253	7	4,260	63.54	15.69
(%)	(0.2)	(1.1)	(3.9)	(12.6)	(22.7)	(25.9)	(19.3)	(8.9)	(4.0)	(0.9)	(0.3)	(0.2)	(100.0)				
14～	3	14	67	204	360	409	256	147	62	22	3	3	1,550	1	1,551	63.27	15.82
(%)	(0.2)	(0.9)	(4.3)	(13.2)	(23.2)	(26.4)	(16.5)	(9.5)	(4.0)	(1.4)	(0.2)	(0.2)	(100.0)				
合計	473	2,452	10,238	26,509	42,172	42,108	27,976	13,028	4,731	1,326	386	200	171,599	232	171,831	60.93	15.63
(%)	(0.3)	(1.4)	(6.0)	(15.4)	(24.6)	(24.5)	(16.3)	(7.6)	(2.8)	(0.8)	(0.2)	(0.1)	(100.0)				
記載なし	6	28	120	302	487	535	346	184	72	22	8	4	2,114	4,264	6,378	62.18	16.44
(%)	(0.3)	(1.3)	(5.7)	(14.3)	(23.0)	(25.3)	(16.4)	(8.7)	(3.4)	(1.0)	(0.4)	(0.2)	(100.0)				
総計	479	2,480	10,358	26,811	42,659	42,643	28,322	13,212	4,803	1,348	394	204	173,713	4,496	178,209	60.94	15.64
(%)	(0.3)	(1.4)	(6.0)	(15.4)	(24.6)	(24.5)	(16.3)	(7.6)	(2.8)	(0.8)	(0.2)	(0.1)	(100.0)				
平均	10.08	10.16	10.39	10.58	10.70	10.79	10.83	10.84	10.83	10.81	10.68	10.48	10.71	10.51	10.71		
標準偏差	1.52	1.46	1.32	1.27	1.23	1.22	1.22	1.24	1.33	1.39	1.39	1.56	1.25	1.37	1.25		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査