

表 2907 透析前BUN(mg/dL) 透析前リン濃度(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

透析前リン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
3未満	13	88	185	219	215	142	69	25	9	1	1		967	1	968	49.28	15.94
(%)	(1.3)	(9.1)	(19.1)	(22.6)	(22.2)	(14.7)	(7.1)	(2.6)	(0.9)	(0.1)	(0.1)		(100.0)				
3.0～	5	55	212	325	336	214	109	40	15	3			1,314		1,314	52.27	14.85
(%)	(0.4)	(4.2)	(16.1)	(24.7)	(25.6)	(16.3)	(8.3)	(3.0)	(1.1)	(0.2)			(100.0)				
3.5～	10	47	302	563	648	482	242	94	26	7			2,421	1	2,422	54.67	14.37
(%)	(0.4)	(1.9)	(12.5)	(23.3)	(26.8)	(19.9)	(10.0)	(3.9)	(1.1)	(0.3)			(100.0)				
4.0～	2	55	295	858	1,113	918	488	164	54	11	1	1	3,960	2	3,962	57.20	13.75
(%)	(0.1)	(1.4)	(7.4)	(21.7)	(28.1)	(23.2)	(12.3)	(4.1)	(1.4)	(0.3)	(0.0)	(0.0)	(100.0)				
4.5～		19	262	870	1,413	1,290	743	278	77	25	7	2	4,986	5	4,991	59.85	13.68
(%)		(0.4)	(5.3)	(17.4)	(28.3)	(25.9)	(14.9)	(5.6)	(1.5)	(0.5)	(0.1)	(0.0)	(100.0)				
5.0～	4	14	179	744	1,494	1,488	993	390	119	23	5	3	5,456	4	5,460	62.01	13.52
(%)	(0.1)	(0.3)	(3.3)	(13.6)	(27.4)	(27.3)	(18.2)	(7.1)	(2.2)	(0.4)	(0.1)	(0.1)	(100.0)				
5.5～	2	10	89	522	1,173	1,397	898	391	141	31	9	1	4,664	4	4,668	63.89	13.45
(%)	(0.0)	(0.2)	(1.9)	(11.2)	(25.2)	(30.0)	(19.3)	(8.4)	(3.0)	(0.7)	(0.2)	(0.0)	(100.0)				
6.0～		2	19	84	193	241	189	89	18	5	2	1	843	1	844	64.64	13.56
(%)		(0.2)	(2.3)	(10.0)	(22.9)	(28.6)	(22.4)	(10.6)	(2.1)	(0.6)	(0.2)	(0.1)	(100.0)				
6.1～		2	47	208	639	897	625	326	109	23	6		2,882	2	2,884	66.33	13.21
(%)		(0.1)	(1.6)	(7.2)	(22.2)	(31.1)	(21.7)	(11.3)	(3.8)	(0.8)	(0.2)		(100.0)				
6.5～		5	23	174	514	688	608	316	112	25	3	5	2,473	1	2,474	67.71	13.86
(%)		(0.2)	(0.9)	(7.0)	(20.8)	(27.8)	(24.6)	(12.8)	(4.5)	(1.0)	(0.1)	(0.2)	(100.0)				
7.0～		2	18	88	268	440	405	226	101	30	7	1	1,586	1	1,587	69.69	14.20
(%)		(0.1)	(1.1)	(5.5)	(16.9)	(27.7)	(25.5)	(14.2)	(6.4)	(1.9)	(0.4)	(0.1)	(100.0)				
7.5～		2	6	45	139	254	248	175	77	25	7		978		978	71.86	14.32
(%)		(0.2)	(0.6)	(4.6)	(14.2)	(26.0)	(25.4)	(17.9)	(7.9)	(2.6)	(0.7)		(100.0)				
8.0～		2	7	24	72	138	167	101	44	17	5	1	578		578	72.45	15.09
(%)		(0.3)	(1.2)	(4.2)	(12.5)	(23.9)	(28.9)	(17.5)	(7.6)	(2.9)	(0.9)	(0.2)	(100.0)				
8.5～	1		1	13	37	69	101	82	43	12	5		364		364	75.10	15.08
(%)	(0.3)		(0.3)	(3.6)	(10.2)	(19.0)	(27.7)	(22.5)	(11.8)	(3.3)	(1.4)		(100.0)				
9.0～	2	1	3	31	41	113	135	116	101	37	16	5	601		601	77.42	17.48
(%)	(0.3)	(0.2)	(0.5)	(5.2)	(6.8)	(18.8)	(22.5)	(19.3)	(16.8)	(6.2)	(2.7)	(0.8)	(100.0)				
合計	39	304	1,648	4,768	8,295	8,771	6,020	2,813	1,046	275	74	20	34,073	22	34,095	62.21	15.08
(%)	(0.1)	(0.9)	(4.8)	(14.0)	(24.3)	(25.7)	(17.7)	(8.3)	(3.1)	(0.8)	(0.2)	(0.1)	(100.0)				
記載なし		5	5	17	23	33	66	18	5	4			176	700	876	67.22	15.92
(%)		(2.8)	(2.8)	(9.7)	(13.1)	(18.8)	(37.5)	(10.2)	(2.8)	(2.3)			(100.0)				
総計	39	309	1,653	4,785	8,318	8,804	6,086	2,831	1,051	279	74	20	34,249	722	34,971	62.23	15.08
(%)	(0.1)	(0.9)	(4.8)	(14.0)	(24.3)	(25.7)	(17.8)	(8.3)	(3.1)	(0.8)	(0.2)	(0.1)	(100.0)				
平均	3.69	3.67	4.24	4.77	5.15	5.48	5.78	6.10	6.48	6.84	7.23	6.95	5.37	5.18	5.37		
標準偏差	2.14	1.29	1.15	1.20	1.20	1.28	1.39	1.52	1.72	1.98	2.05	2.00	1.42	0.99	1.42		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査