

表 2919 BUN(mg/dL) 透析歴(年)別 (腹膜透析患者)

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
2年未満	11	86	286	541	621	471	222	107	40	23	2	3	2,413	1,187	3,600	55.04	16.44
(%)	(0.5)	(3.6)	(11.9)	(22.4)	(25.7)	(19.5)	(9.2)	(4.4)	(1.7)	(1.0)	(0.1)	(0.1)	(100.0)				
2年～	10	49	222	484	587	392	195	89	32	14	4	2	2,080	1,121	3,201	55.38	15.82
(%)	(0.5)	(2.4)	(10.7)	(23.3)	(28.2)	(18.8)	(9.4)	(4.3)	(1.5)	(0.7)	(0.2)	(0.1)	(100.0)				
5年～	5	38	123	287	291	231	110	37	12	3	1	2	1,140	517	1,657	54.33	14.88
(%)	(0.4)	(3.3)	(10.8)	(25.2)	(25.5)	(20.3)	(9.6)	(3.2)	(1.1)	(0.3)	(0.1)	(0.2)	(100.0)				
10年～	1	5	27	44	82	42	19	9	4				233	107	340	54.49	14.18
(%)	(0.4)	(2.1)	(11.6)	(18.9)	(35.2)	(18.0)	(8.2)	(3.9)	(1.7)				(100.0)				
15年～		3	9	23	26	9	2		1				73	24	97	49.90	11.36
(%)		(4.1)	(12.3)	(31.5)	(35.6)	(12.3)	(2.7)		(1.4)				(100.0)				
20年～	1	1	1	7	3	2		1					16	4	20	46.44	16.03
(%)	(6.3)	(6.3)	(6.3)	(43.8)	(18.8)	(12.5)		(6.3)					(100.0)				
25年～		1		3	1	2		1					8		8	55.13	17.96
(%)		(12.5)		(37.5)	(12.5)	(25.0)		(12.5)					(100.0)				
30年～	1		1	2	2	1	1						8	2	10	48.00	21.41
(%)	(12.5)		(12.5)	(25.0)	(25.0)	(12.5)	(12.5)						(100.0)				
35年～			2	1									3		3	37.00	7.00
(%)			(66.7)	(33.3)									(100.0)				
40年～																	
(%)																	
合計	29	183	671	1,392	1,613	1,150	549	244	89	40	7	7	5,974	2,962	8,936	54.90	15.82
(%)	(0.5)	(3.1)	(11.2)	(23.3)	(27.0)	(19.3)	(9.2)	(4.1)	(1.5)	(0.7)	(0.1)	(0.1)	(100.0)				
不明			1					1					2	3	5	63.50	36.06
(%)			(50.0)					(50.0)					(100.0)				
記載なし																	
(%)																	
総計	29	183	672	1,392	1,613	1,150	549	245	89	40	7	7	5,976	2,965	8,941	54.90	15.82
(%)	(0.5)	(3.1)	(11.2)	(23.3)	(27.0)	(19.2)	(9.2)	(4.1)	(1.5)	(0.7)	(0.1)	(0.1)	(100.0)				
平均	4.41	3.14	3.14	3.35	3.28	3.03	2.92	2.82	2.71	1.55	2.43	3.14	3.16	2.98	3.10		
標準偏差	7.21	4.14	4.10	3.96	3.67	3.39	3.17	3.42	3.03	1.66	2.37	2.34	3.71	3.25	3.56		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査