

表 2920 BUN(mg/dL) 原疾患別（腹膜透析患者）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	8	46	188	454	582	422	231	94	39	11	2	1	2,078	981	3,059	56.43	15.23
(%)	(0.4)	(2.2)	(9.0)	(21.8)	(28.0)	(20.3)	(11.1)	(4.5)	(1.9)	(0.5)	(0.1)	(0.0)	(100.0)				
慢性腎盂腎炎		4	4	22	13	13	8	2	1	1			68	25	93	54.66	16.27
(%)		(5.9)	(5.9)	(32.4)	(19.1)	(19.1)	(11.8)	(2.9)	(1.5)	(1.5)			(100.0)				
急速進行性糸球体腎炎	1	1	3	10	16	15	1		1	1			49	25	74	54.69	15.34
(%)	(2.0)	(2.0)	(6.1)	(20.4)	(32.7)	(30.6)	(2.0)		(2.0)	(2.0)			(100.0)				
妊娠腎/妊娠中毒症			2	8	7	8							25	7	32	53.68	9.52
(%)			(8.0)	(32.0)	(28.0)	(32.0)							(100.0)				
その他の分類不能の腎炎			6	13	20	11	7	2					59	26	85	55.12	12.41
(%)			(10.2)	(22.0)	(33.9)	(18.6)	(11.9)	(3.4)					(100.0)				
多発性嚢胞腎		5	13	43	34	29	15	11	2	3		1	156	87	243	57.18	17.25
(%)		(3.2)	(8.3)	(27.6)	(21.8)	(18.6)	(9.6)	(7.1)	(1.3)	(1.9)		(0.6)	(100.0)				
腎硬化症	2	21	73	169	209	147	73	33	11	5		2	745	338	1,083	55.81	16.45
(%)	(0.3)	(2.8)	(9.8)	(22.7)	(28.1)	(19.7)	(9.8)	(4.4)	(1.5)	(0.7)		(0.3)	(100.0)				
悪性高血圧	1	1	9	9	21	14	8	3	2		1		69	38	107	57.14	16.96
(%)	(1.4)	(1.4)	(13.0)	(13.0)	(30.4)	(20.3)	(11.6)	(4.3)	(2.9)		(1.4)		(100.0)				
糖尿病性腎症	10	69	265	445	476	299	130	65	22	11	2	1	1,795	971	2,766	52.79	15.34
(%)	(0.6)	(3.8)	(14.8)	(24.8)	(26.5)	(16.7)	(7.2)	(3.6)	(1.2)	(0.6)	(0.1)	(0.1)	(100.0)				
SLE腎炎		4	7	9	9	8	3	4	1				45	17	62	53.47	17.84
(%)		(8.9)	(15.6)	(20.0)	(20.0)	(17.8)	(6.7)	(8.9)	(2.2)				(100.0)				
アミロイド腎				3		1							4	5	9	49.75	7.80
(%)				(75.0)		(25.0)							(100.0)				
痛風腎			2	2	7	10	2	1	1				25	8	33	60.16	13.86
(%)			(8.0)	(8.0)	(28.0)	(40.0)	(8.0)	(4.0)	(4.0)				(100.0)				
先天性代謝異常による腎不全		2	3		2	3	2	1			1		14	4	18	57.50	24.68
(%)		(14.3)	(21.4)		(14.3)	(21.4)	(14.3)	(7.1)			(7.1)		(100.0)				
腎・尿路結核					1								1		1	51.00	
(%)					(100.0)								(100.0)				
腎・尿路結石		1	1	2	2	3	1						10	6	16	52.30	14.45
(%)		(10.0)	(10.0)	(20.0)	(20.0)	(30.0)	(10.0)						(100.0)				
腎・尿路腫瘍			3		3	3							9	5	14	51.44	13.49
(%)			(33.3)		(33.3)	(33.3)							(100.0)				
閉塞性尿路障害			2	4	4	2	3			1			16	6	22	57.69	19.02
(%)			(12.5)	(25.0)	(25.0)	(12.5)	(18.8)			(6.3)			(100.0)				
骨髄腫		1	1	1	2								5	3	8	41.60	12.01
(%)		(20.0)	(20.0)	(20.0)	(40.0)								(100.0)				
腎形成不全			5	10	24	12		3		1			55	21	76	55.40	13.24
(%)			(9.1)	(18.2)	(43.6)	(21.8)		(5.5)		(1.8)			(100.0)				
不明	4	21	63	135	137	114	46	16	6	3	1	2	548	292	840	54.08	17.33
(%)	(0.7)	(3.8)	(11.5)	(24.6)	(25.0)	(20.8)	(8.4)	(2.9)	(1.1)	(0.5)	(0.2)	(0.4)	(100.0)				
再導入			1	10	11	10		3	1				36	15	51	56.56	14.27
(%)			(2.8)	(27.8)	(30.6)	(27.8)		(8.3)	(2.8)				(100.0)				
その他	3	7	21	43	33	26	19	7	2	3			164	85	249	53.94	17.56
(%)	(1.8)	(4.3)	(12.8)	(26.2)	(20.1)	(15.9)	(11.6)	(4.3)	(1.2)	(1.8)			(100.0)				
合計	29	183	672	1,392	1,613	1,150	549	245	89	40	7	7	5,976	2,965	8,941	54.90	15.82
(%)	(0.5)	(3.1)	(11.2)	(23.3)	(27.0)	(19.2)	(9.2)	(4.1)	(1.5)	(0.7)	(0.1)	(0.1)	(100.0)				
記載なし																	
(%)																	
総計	29	183	672	1,392	1,613	1,150	549	245	89	40	7	7	5,976	2,965	8,941	54.90	15.82
(%)	(0.5)	(3.1)	(11.2)	(23.3)	(27.0)	(19.2)	(9.2)	(4.1)	(1.5)	(0.7)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査