

表 2924 BUN(mg/dL) アルブミン濃度(g/dL)別（腹膜透析患者）

アルブミン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
1.5未満	1	2	3	1	1	2	1		3				14	1	15	53.57	26.83
(%)	(7.1)	(14.3)	(21.4)	(7.1)	(7.1)	(14.3)	(7.1)		(21.4)				(100.0)				
1.5～	3	10	19	20	10	5	2	1					70	1	71	41.94	14.07
(%)	(4.3)	(14.3)	(27.1)	(28.6)	(14.3)	(7.1)	(2.9)	(1.4)					(100.0)				
2.0～	5	33	55	72	49	40	16	12	4	6	1	1	294	4	298	50.10	22.06
(%)	(1.7)	(11.2)	(18.7)	(24.5)	(16.7)	(13.6)	(5.4)	(4.1)	(1.4)	(2.0)	(0.3)	(0.3)	(100.0)				
2.5～	7	46	173	266	249	139	89	31	19	5	1	1	1,026	11	1,037	52.06	16.86
(%)	(0.7)	(4.5)	(16.9)	(25.9)	(24.3)	(13.5)	(8.7)	(3.0)	(1.9)	(0.5)	(0.1)	(0.1)	(100.0)				
3.0～	5	45	224	508	594	433	197	93	27	13	1	2	2,142	16	2,158	55.42	14.84
(%)	(0.2)	(2.1)	(10.5)	(23.7)	(27.7)	(20.2)	(9.2)	(4.3)	(1.3)	(0.6)	(0.0)	(0.1)	(100.0)				
3.5～	5	37	155	422	554	400	187	85	31	12	4	2	1,894	12	1,906	56.54	14.80
(%)	(0.3)	(2.0)	(8.2)	(22.3)	(29.3)	(21.1)	(9.9)	(4.5)	(1.6)	(0.6)	(0.2)	(0.1)	(100.0)				
4.0～	3	7	27	75	133	110	46	18	5	3			427	4	431	57.47	14.29
(%)	(0.7)	(1.6)	(6.3)	(17.6)	(31.1)	(25.8)	(10.8)	(4.2)	(1.2)	(0.7)			(100.0)				
4.5～			2	5	4	5	2	3					21		21	59.62	15.02
(%)			(9.5)	(23.8)	(19.0)	(23.8)	(9.5)	(14.3)					(100.0)				
5.0～			1				1						2		2	57.50	30.41
(%)			(50.0)				(50.0)						(100.0)				
合計	29	180	659	1,369	1,594	1,134	541	243	89	39	7	6	5,890	49	5,939	54.93	15.81
(%)	(0.5)	(3.1)	(11.2)	(23.2)	(27.1)	(19.3)	(9.2)	(4.1)	(1.5)	(0.7)	(0.1)	(0.1)	(100.0)				
記載なし		3	13	23	19	16	8	2		1		1	86	2,916	3,002	53.13	16.97
(%)		(3.5)	(15.1)	(26.7)	(22.1)	(18.6)	(9.3)	(2.3)		(1.2)		(1.2)	(100.0)				
総計	29	183	672	1,392	1,613	1,150	549	245	89	40	7	7	5,976	2,965	8,941	54.90	15.82
(%)	(0.5)	(3.1)	(11.2)	(23.3)	(27.0)	(19.2)	(9.2)	(4.1)	(1.5)	(0.7)	(0.1)	(0.1)	(100.0)				
平均	2.86	2.94	3.09	3.24	3.34	3.37	3.36	3.33	3.22	3.21	3.33	3.12	3.28	3.13	3.28		
標準偏差	0.79	0.64	0.57	0.51	0.47	0.48	0.49	0.49	0.60	0.56	0.68	0.63	0.52	0.60	0.52		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査