

表 2933 BUN(mg/dL) Non-HDL-C濃度(mg/dL)別（腹膜透析患者）

Non-HDL-C濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
50未満		1	2	4	6	4	1						18		18	53.17	13.54
(%)		(5.6)	(11.1)	(22.2)	(33.3)	(22.2)	(5.6)						(100.0)				
50～	1	4	9	22	33	19	15	5	3	1			112	2	114	56.51	16.17
(%)	(0.9)	(3.6)	(8.0)	(19.6)	(29.5)	(17.0)	(13.4)	(4.5)	(2.7)	(0.9)			(100.0)				
70～	3	11	32	66	93	75	34	14	10	4		1	343	8	351	56.73	16.53
(%)	(0.9)	(3.2)	(9.3)	(19.2)	(27.1)	(21.9)	(9.9)	(4.1)	(2.9)	(1.2)		(0.3)	(100.0)				
90～	1	19	85	143	184	119	65	27	8	1			652	8	660	54.28	14.60
(%)	(0.2)	(2.9)	(13.0)	(21.9)	(28.2)	(18.3)	(10.0)	(4.1)	(1.2)	(0.2)			(100.0)				
110～	1	21	80	195	187	148	92	42	9	8		1	784	13	797	56.01	15.57
(%)	(0.1)	(2.7)	(10.2)	(24.9)	(23.9)	(18.9)	(11.7)	(5.4)	(1.1)	(1.0)		(0.1)	(100.0)				
130～	2	14	68	157	178	132	64	21	10	2	1		649	13	662	55.23	14.48
(%)	(0.3)	(2.2)	(10.5)	(24.2)	(27.4)	(20.3)	(9.9)	(3.2)	(1.5)	(0.3)	(0.2)		(100.0)				
150～	1	15	43	117	114	80	33	21	4	3	1		432	4	436	54.18	15.41
(%)	(0.2)	(3.5)	(10.0)	(27.1)	(26.4)	(18.5)	(7.6)	(4.9)	(0.9)	(0.7)	(0.2)		(100.0)				
170～	3	4	35	61	72	52	22	7	2	3	1		262	1	263	54.14	15.20
(%)	(1.1)	(1.5)	(13.4)	(23.3)	(27.5)	(19.8)	(8.4)	(2.7)	(0.8)	(1.1)	(0.4)		(100.0)				
190～		3	16	29	34	25	5	4	1	3			120		120	54.01	15.60
(%)		(2.5)	(13.3)	(24.2)	(28.3)	(20.8)	(4.2)	(3.3)	(0.8)	(2.5)			(100.0)				
210～	1	11	11	14	27	14	5	2	1	1			87	3	90	50.80	16.58
(%)	(1.1)	(12.6)	(12.6)	(16.1)	(31.0)	(16.1)	(5.7)	(2.3)	(1.1)	(1.1)			(100.0)				
合計	13	103	381	808	928	668	336	143	48	26	3	2	3,459	52	3,511	55.04	15.30
(%)	(0.4)	(3.0)	(11.0)	(23.4)	(26.8)	(19.3)	(9.7)	(4.1)	(1.4)	(0.8)	(0.1)	(0.1)	(100.0)				
記載なし	16	80	291	584	685	482	213	102	41	14	4	5	2,517	2,913	5,430	54.72	16.51
(%)	(0.6)	(3.2)	(11.6)	(23.2)	(27.2)	(19.1)	(8.5)	(4.1)	(1.6)	(0.6)	(0.2)	(0.2)	(100.0)				
総計	29	183	672	1,392	1,613	1,150	549	245	89	40	7	7	5,976	2,965	8,941	54.90	15.82
(%)	(0.5)	(3.1)	(11.2)	(23.3)	(27.0)	(19.2)	(9.2)	(4.1)	(1.5)	(0.7)	(0.1)	(0.1)	(100.0)				
平均	133.92	133.49	130.58	130.03	128.80	128.37	122.83	124.50	118.81	136.08	161.33	100.50	128.53	124.69	128.47		
標準偏差	51.57	44.33	39.49	36.44	39.66	37.24	35.21	33.45	34.42	43.30	21.55	20.51	37.98	40.46	38.01		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査