

表 2940 透析前クレアチニン濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	2未満	2～	4～	6～	8～	10～	12～	14～	16～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	13	339	1,743	5,897	13,700	16,975	11,809	5,440	1,991	57,907	1,540	59,447	10.85	2.77
(%)	(0.0)	(0.6)	(3.0)	(10.2)	(23.7)	(29.3)	(20.4)	(9.4)	(3.4)	(100.0)				
慢性腎盂腎炎	4	12	73	244	486	520	271	125	47	1,782	35	1,817	10.28	2.73
(%)	(0.2)	(0.7)	(4.1)	(13.7)	(27.3)	(29.2)	(15.2)	(7.0)	(2.6)	(100.0)				
急速進行性糸球体腎炎	5	31	94	216	330	296	170	70	35	1,247	34	1,281	9.68	3.06
(%)	(0.4)	(2.5)	(7.5)	(17.3)	(26.5)	(23.7)	(13.6)	(5.6)	(2.8)	(100.0)				
妊娠腎/妊娠中毒症		4	22	111	313	357	145	23	2	977	22	999	10.10	2.07
(%)		(0.4)	(2.3)	(11.4)	(32.0)	(36.5)	(14.8)	(2.4)	(0.2)	(100.0)				
その他の分類不能の腎炎		13	35	74	134	205	157	87	40	745	18	763	11.05	3.15
(%)		(1.7)	(4.7)	(9.9)	(18.0)	(27.5)	(21.1)	(11.7)	(5.4)	(100.0)				
多発性嚢胞腎	1	18	149	672	1,713	2,071	1,292	519	186	6,621	163	6,784	10.75	2.61
(%)	(0.0)	(0.3)	(2.3)	(10.1)	(25.9)	(31.3)	(19.5)	(7.8)	(2.8)	(100.0)				
腎硬化症	7	156	762	2,403	4,162	3,686	2,075	815	263	14,329	348	14,677	9.92	2.76
(%)	(0.0)	(1.1)	(5.3)	(16.8)	(29.0)	(25.7)	(14.5)	(5.7)	(1.8)	(100.0)				
悪性高血圧	1	14	56	167	313	361	269	164	69	1,414	51	1,465	10.87	3.05
(%)	(0.1)	(1.0)	(4.0)	(11.8)	(22.1)	(25.5)	(19.0)	(11.6)	(4.9)	(100.0)				
糖尿病性腎症	19	753	3,573	10,893	19,058	17,538	9,524	3,038	758	65,154	1,696	66,850	9.83	2.65
(%)	(0.0)	(1.2)	(5.5)	(16.7)	(29.3)	(26.9)	(14.6)	(4.7)	(1.2)	(100.0)				
SLE腎炎	1	12	72	171	334	345	183	43	20	1,181	34	1,215	9.92	2.62
(%)	(0.1)	(1.0)	(6.1)	(14.5)	(28.3)	(29.2)	(15.5)	(3.6)	(1.7)	(100.0)				
アミロイド腎		7	23	57	66	35	24	4	1	217	13	230	8.70	2.70
(%)		(3.2)	(10.6)	(26.3)	(30.4)	(16.1)	(11.1)	(1.8)	(0.5)	(100.0)				
痛風腎		1	11	59	152	190	167	70	39	689	16	705	11.31	2.75
(%)		(0.1)	(1.6)	(8.6)	(22.1)	(27.6)	(24.2)	(10.2)	(5.7)	(100.0)				
先天性代謝異常による腎不全		1	4	13	24	37	40	20	12	151	3	154	11.66	3.09
(%)		(0.7)	(2.6)	(8.6)	(15.9)	(24.5)	(26.5)	(13.2)	(7.9)	(100.0)				
腎・尿路結核		3	10	16	49	37	17	1		133	2	135	9.41	2.38
(%)		(2.3)	(7.5)	(12.0)	(36.8)	(27.8)	(12.8)	(0.8)		(100.0)				
腎・尿路結石		2	13	41	89	101	65	27	8	346	8	354	10.51	2.69
(%)		(0.6)	(3.8)	(11.8)	(25.7)	(29.2)	(18.8)	(7.8)	(2.3)	(100.0)				
腎・尿路腫瘍	1	3	27	49	107	136	85	35	17	460	13	473	10.56	2.85
(%)	(0.2)	(0.7)	(5.9)	(10.7)	(23.3)	(29.6)	(18.5)	(7.6)	(3.7)	(100.0)				
閉塞性尿路障害		17	19	82	86	80	66	34	19	403	11	414	10.05	3.34
(%)		(4.2)	(4.7)	(20.3)	(21.3)	(19.9)	(16.4)	(8.4)	(4.7)	(100.0)				
骨髄腫		2	8	21	24	15	9	4		83	5	88	9.01	2.73
(%)		(2.4)	(9.6)	(25.3)	(28.9)	(18.1)	(10.8)	(4.8)		(100.0)				
腎形成不全			4	18	40	69	110	81	32	354	5	359	12.61	2.71
(%)			(1.1)	(5.1)	(11.3)	(19.5)	(31.1)	(22.9)	(9.0)	(100.0)				
不明	11	189	854	2,144	3,794	3,693	2,321	951	388	14,345	485	14,830	10.12	2.94
(%)	(0.1)	(1.3)	(6.0)	(14.9)	(26.4)	(25.7)	(16.2)	(6.6)	(2.7)	(100.0)				
再導入		5	27	90	189	323	311	166	49	1,160	22	1,182	11.52	2.76
(%)		(0.4)	(2.3)	(7.8)	(16.3)	(27.8)	(26.8)	(14.3)	(4.2)	(100.0)				
その他	13	74	216	568	952	964	644	319	124	3,874	113	3,987	10.24	3.08
(%)	(0.3)	(1.9)	(5.6)	(14.7)	(24.6)	(24.9)	(16.6)	(8.2)	(3.2)	(100.0)				
合計	76	1,656	7,795	24,006	46,115	48,034	29,754	12,036	4,100	173,572	4,637	178,209	10.29	2.79
(%)	(0.0)	(1.0)	(4.5)	(13.8)	(26.6)	(27.7)	(17.1)	(6.9)	(2.4)	(100.0)				
記載なし														
(%)														
総計	76	1,656	7,795	24,006	46,115	48,034	29,754	12,036	4,100	173,572	4,637	178,209	10.29	2.79
(%)	(0.0)	(1.0)	(4.5)	(13.8)	(26.6)	(27.7)	(17.1)	(6.9)	(2.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査