

表 2945 透析前クレアチニン濃度(mg/dL) nPCR(g/kg/day)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

nPCR	2未満	2～	4～	6～	8～	10～	12～	14～	16～	合計	記載なし	総計	平均	標準偏差
0.4未満	11	50	117	82	46	33	17	6	7	369		369	6.93	4.16
(%)	(3.0)	(13.6)	(31.7)	(22.2)	(12.5)	(8.9)	(4.6)	(1.6)	(1.9)	(100.0)				
0.4～	32	534	1,618	2,691	2,174	1,005	282	63	19	8,418		8,418	7.58	2.51
(%)	(0.4)	(6.3)	(19.2)	(32.0)	(25.8)	(11.9)	(3.3)	(0.7)	(0.2)	(100.0)				
0.6～	18	594	3,195	10,071	15,920	12,652	6,189	1,840	477	50,956		50,956	9.52	2.61
(%)	(0.0)	(1.2)	(6.3)	(19.8)	(31.2)	(24.8)	(12.1)	(3.6)	(0.9)	(100.0)				
0.8～	5	222	1,594	7,332	18,377	21,502	13,943	5,720	1,816	70,511		70,511	10.74	2.57
(%)	(0.0)	(0.3)	(2.3)	(10.4)	(26.1)	(30.5)	(19.8)	(8.1)	(2.6)	(100.0)				
1.0～		44	405	1,909	6,050	8,775	6,686	3,159	1,220	28,248		28,248	11.35	2.60
(%)		(0.2)	(1.4)	(6.8)	(21.4)	(31.1)	(23.7)	(11.2)	(4.3)	(100.0)				
1.2～		18	87	293	1,007	1,539	1,137	574	261	4,916		4,916	11.45	2.69
(%)		(0.4)	(1.8)	(6.0)	(20.5)	(31.3)	(23.1)	(11.7)	(5.3)	(100.0)				
1.4～		3	9	28	96	168	127	81	43	555		555	11.97	3.42
(%)		(0.5)	(1.6)	(5.0)	(17.3)	(30.3)	(22.9)	(14.6)	(7.7)	(100.0)				
1.6～		1	4	13	12	23	11	11	4	79		79	10.69	3.21
(%)		(1.3)	(5.1)	(16.5)	(15.2)	(29.1)	(13.9)	(13.9)	(5.1)	(100.0)				
1.8～			1	2	12	3	5	1	1	25		25	10.36	2.65
(%)			(4.0)	(8.0)	(48.0)	(12.0)	(20.0)	(4.0)	(4.0)	(100.0)				
2.0～		1		3	6	15	4	6	3	38		38	11.78	3.88
(%)		(2.6)		(7.9)	(15.8)	(39.5)	(10.5)	(15.8)	(7.9)	(100.0)				
合計	66	1,467	7,030	22,424	43,700	45,715	28,401	11,461	3,851	164,115		164,115	10.32	2.77
(%)	(0.0)	(0.9)	(4.3)	(13.7)	(26.6)	(27.9)	(17.3)	(7.0)	(2.3)	(100.0)				
記載なし	10	189	765	1,582	2,415	2,319	1,353	575	249	9,457	4,637	14,094	9.81	3.05
(%)	(0.1)	(2.0)	(8.1)	(16.7)	(25.5)	(24.5)	(14.3)	(6.1)	(2.6)	(100.0)				
総計	76	1,656	7,795	24,006	46,115	48,034	29,754	12,036	4,100	173,572	4,637	178,209	10.29	2.79
(%)	(0.0)	(1.0)	(4.5)	(13.8)	(26.6)	(27.7)	(17.1)	(6.9)	(2.4)	(100.0)				
平均	0.55	0.66	0.72	0.78	0.84	0.89	0.91	0.94	0.97	0.86		0.86		
標準偏差	0.15	0.19	0.18	0.17	0.17	0.16	0.16	0.16	0.17	0.18		0.18		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査