

表 2948 透析前クレアチニン濃度(mg/dL) 透析前CRP濃度(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前CRP濃度	2未満	2～	4～	6～	8～	10～	12～	14～	16～	合計	記載なし	総計	平均	標準偏差
0.2未満	38	437	2,311	9,379	21,094	24,273	15,593	6,241	2,065	81,431	75	81,506	10.61	2.65
(%)	(0.0)	(0.5)	(2.8)	(11.5)	(25.9)	(29.8)	(19.1)	(7.7)	(2.5)	(100.0)				
0.2～	9	235	1,277	4,045	7,522	7,615	4,645	1,854	616	27,818	22	27,840	10.22	2.75
(%)	(0.0)	(0.8)	(4.6)	(14.5)	(27.0)	(27.4)	(16.7)	(6.7)	(2.2)	(100.0)				
0.5～	3	179	879	2,438	4,089	3,538	2,025	798	277	14,226	20	14,246	9.86	2.84
(%)	(0.0)	(1.3)	(6.2)	(17.1)	(28.7)	(24.9)	(14.2)	(5.6)	(1.9)	(100.0)				
1.0～	7	189	784	1,876	2,563	2,122	1,095	433	161	9,230	15	9,245	9.45	2.92
(%)	(0.1)	(2.0)	(8.5)	(20.3)	(27.8)	(23.0)	(11.9)	(4.7)	(1.7)	(100.0)				
2.0～	6	160	668	1,220	1,644	1,162	521	184	70	5,635	8	5,643	8.97	2.89
(%)	(0.1)	(2.8)	(11.9)	(21.7)	(29.2)	(20.6)	(9.2)	(3.3)	(1.2)	(100.0)				
4.0～		81	295	467	470	375	202	60	18	1,968	8	1,976	8.73	3.34
(%)		(4.1)	(15.0)	(23.7)	(23.9)	(19.1)	(10.3)	(3.0)	(0.9)	(100.0)				
6.0～		35	155	240	237	160	76	28	11	942	7	949	8.52	2.87
(%)		(3.7)	(16.5)	(25.5)	(25.2)	(17.0)	(8.1)	(3.0)	(1.2)	(100.0)				
8.0～		30	80	129	125	85	56	20	5	530	2	532	8.58	3.23
(%)		(5.7)	(15.1)	(24.3)	(23.6)	(16.0)	(10.6)	(3.8)	(0.9)	(100.0)				
10.0～	1	28	107	143	157	124	65	21	5	651		651	8.58	3.05
(%)	(0.2)	(4.3)	(16.4)	(22.0)	(24.1)	(19.0)	(10.0)	(3.2)	(0.8)	(100.0)				
15.0～		21	72	107	128	65	34	13	5	445	4	449	8.43	2.88
(%)		(4.7)	(16.2)	(24.0)	(28.8)	(14.6)	(7.6)	(2.9)	(1.1)	(100.0)				
合計	64	1,395	6,628	20,044	38,029	39,519	24,312	9,652	3,233	142,876	161	143,037	10.26	2.78
(%)	(0.0)	(1.0)	(4.6)	(14.0)	(26.6)	(27.7)	(17.0)	(6.8)	(2.3)	(100.0)				
記載なし	12	261	1,167	3,962	8,086	8,515	5,442	2,384	867	30,696	4,476	35,172	10.47	2.80
(%)	(0.0)	(0.9)	(3.8)	(12.9)	(26.3)	(27.7)	(17.7)	(7.8)	(2.8)	(100.0)				
総計	76	1,656	7,795	24,006	46,115	48,034	29,754	12,036	4,100	173,572	4,637	178,209	10.29	2.79
(%)	(0.0)	(1.0)	(4.5)	(13.8)	(26.6)	(27.7)	(17.1)	(6.9)	(2.4)	(100.0)				
平均	0.78	1.89	1.55	0.95	0.65	0.49	0.43	0.41	0.40	0.64	1.61	0.64		
標準偏差	1.96	3.52	3.12	2.52	2.02	1.59	1.46	1.34	1.15	1.96	3.51	1.97		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査