

表 2951 透析前クレアチニン濃度(mg/dL) PTH値(pg/mL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上) PTH測定法:intact-PTH

PTH値	2未満	2～	4～	6～	8～	10～	12～	14～	16～	合計	記載なし	総計	平均	標準偏差
20未満	4	150	663	1,261	2,017	1,888	1,018	360	107	7,468	9	7,477	9.60	2.88
(%)	(0.1)	(2.0)	(8.9)	(16.9)	(27.0)	(25.3)	(13.6)	(4.8)	(1.4)	(100.0)				
20～	15	150	782	1,807	3,014	2,760	1,509	510	160	10,707	17	10,724	9.73	2.82
(%)	(0.1)	(1.4)	(7.3)	(16.9)	(28.1)	(25.8)	(14.1)	(4.8)	(1.5)	(100.0)				
40～	11	199	697	1,867	3,152	3,029	1,705	631	192	11,483	19	11,502	9.87	2.82
(%)	(0.1)	(1.7)	(6.1)	(16.3)	(27.4)	(26.4)	(14.8)	(5.5)	(1.7)	(100.0)				
60～	7	140	652	1,849	3,386	3,375	1,876	659	231	12,175	15	12,190	10.01	2.73
(%)	(0.1)	(1.1)	(5.4)	(15.2)	(27.8)	(27.7)	(15.4)	(5.4)	(1.9)	(100.0)				
80～	5	122	585	1,870	3,440	3,437	1,994	727	230	12,410	26	12,436	10.11	2.73
(%)	(0.0)	(1.0)	(4.7)	(15.1)	(27.7)	(27.7)	(16.1)	(5.9)	(1.9)	(100.0)				
100～	2	123	503	1,657	3,367	3,338	2,045	809	269	12,113	25	12,138	10.27	2.75
(%)	(0.0)	(1.0)	(4.2)	(13.7)	(27.8)	(27.6)	(16.9)	(6.7)	(2.2)	(100.0)				
120～	7	92	434	1,614	3,063	3,216	2,025	786	253	11,490	10	11,500	10.32	2.71
(%)	(0.1)	(0.8)	(3.8)	(14.0)	(26.7)	(28.0)	(17.6)	(6.8)	(2.2)	(100.0)				
140～	1	78	375	1,334	2,868	2,981	1,864	759	242	10,502	17	10,519	10.40	2.66
(%)	(0.0)	(0.7)	(3.6)	(12.7)	(27.3)	(28.4)	(17.7)	(7.2)	(2.3)	(100.0)				
160～	4	64	298	1,252	2,428	2,784	1,781	694	233	9,538	23	9,561	10.50	2.67
(%)	(0.0)	(0.7)	(3.1)	(13.1)	(25.5)	(29.2)	(18.7)	(7.3)	(2.4)	(100.0)				
180～	4	111	643	2,586	5,502	6,073	4,016	1,724	619	21,278	26	21,304	10.62	2.73
(%)	(0.0)	(0.5)	(3.0)	(12.2)	(25.9)	(28.5)	(18.9)	(8.1)	(2.9)	(100.0)				
240～	2	6	27	65	87	46	22	3	258	1	259	10.60	2.47	
(%)		(0.8)	(2.3)	(10.5)	(25.2)	(33.7)	(17.8)	(8.5)	(1.2)	(100.0)				
241～	48	341	1,373	3,008	3,539	2,303	1,067	351	12,030	12	12,042	10.73	2.74	
(%)		(0.4)	(2.8)	(11.4)	(25.0)	(29.4)	(19.1)	(8.9)	(2.9)	(100.0)				
300～	20	159	687	1,734	2,035	1,432	601	258	6,926	11	6,937	10.90	2.69	
(%)		(0.3)	(2.3)	(9.9)	(25.0)	(29.4)	(20.7)	(8.7)	(3.7)	(100.0)				
360～	13	80	386	991	1,111	807	399	140	3,927	3	3,930	10.96	2.69	
(%)		(0.3)	(2.0)	(9.8)	(25.2)	(28.3)	(20.6)	(10.2)	(3.6)	(100.0)				
420～	5	51	248	561	643	489	252	89	2,338	9	2,347	10.99	2.76	
(%)		(0.2)	(2.2)	(10.6)	(24.0)	(27.5)	(20.9)	(10.8)	(3.8)	(100.0)				
480～	4	17	68	119	166	124	53	23	574	2	576	10.93	2.85	
(%)		(0.7)	(3.0)	(11.8)	(20.7)	(28.9)	(21.6)	(9.2)	(4.0)	(100.0)				
500～	1	18	92	217	273	184	108	41	934		934	11.09	2.77	
(%)		(0.1)	(1.9)	(9.9)	(23.2)	(29.2)	(19.7)	(11.6)	(4.4)	(100.0)				
540～	1	24	107	225	294	225	106	44	1,026		1,026	11.09	2.77	
(%)		(0.1)	(2.3)	(10.4)	(21.9)	(28.7)	(21.9)	(10.3)	(4.3)	(100.0)				
600～	1	33	180	418	450	354	179	82	1,697	3	1,700	11.09	2.76	
(%)		(0.1)	(1.9)	(10.6)	(24.6)	(26.5)	(20.9)	(10.5)	(4.8)	(100.0)				
800～	6	51	159	324	363	290	142	67	1,402	3	1,405	10.91	2.96	
(%)		(0.4)	(3.6)	(11.3)	(23.1)	(25.9)	(20.7)	(10.1)	(4.8)	(100.0)				
合計	60	1,330	6,412	20,424	39,899	41,842	26,087	10,588	3,634	150,276	231	150,507	10.33	2.77
(%)	(0.0)	(0.9)	(4.3)	(13.6)	(26.6)	(27.8)	(17.4)	(7.0)	(2.4)	(100.0)				
記載なし	3	65	215	448	652	608	318	134	41	2,484	614	3,098	9.57	3.04
(%)	(0.1)	(2.6)	(8.7)	(18.0)	(26.2)	(24.5)	(12.8)	(5.4)	(1.7)	(100.0)				
総計	63	1,395	6,627	20,872	40,551	42,450	26,405	10,722	3,675	152,760	845	153,605	10.32	2.78
(%)	(0.0)	(0.9)	(4.3)	(13.7)	(26.5)	(27.8)	(17.3)	(7.0)	(2.4)	(100.0)				
平均	78.00	110.27	129.98	152.49	164.06	172.59	184.93	200.07	215.83	170.31	170.16	170.31		
標準偏差	55.48	128.79	167.04	160.51	159.15	161.20	167.38	179.24	201.34	164.97	163.96	164.97		

数値下のかつこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査