

表 2953 透析前クレアチニン濃度(mg/dL) 透析前ヘモグロビン濃度(g/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前ヘモグロビン 濃度	2未満	2～	4～	6～	8～	10～	12～	14～	16～	合計	記載なし	総計	平均	標準偏差
5未満		5	16	36	67	46	30	6	4	210		210	9.62	3.02
(%)		(2.4)	(7.6)	(17.1)	(31.9)	(21.9)	(14.3)	(2.9)	(1.9)	(100.0)				
5～		9	24	57	75	51	30	4	1	251		251	8.95	2.70
(%)		(3.6)	(9.6)	(22.7)	(29.9)	(20.3)	(12.0)	(1.6)	(0.4)	(100.0)				
6～		39	87	170	164	106	37	8	4	615	2	617	8.19	2.78
(%)		(6.3)	(14.1)	(27.6)	(26.7)	(17.2)	(6.0)	(1.3)	(0.7)	(100.0)				
7～	3	95	370	635	699	418	160	42	16	2,438	7	2,445	8.39	2.81
(%)	(0.1)	(3.9)	(15.2)	(26.0)	(28.7)	(17.1)	(6.6)	(1.7)	(0.7)	(100.0)				
8～	6	184	873	1,978	2,541	1,792	812	223	72	8,481	16	8,497	8.97	2.70
(%)	(0.1)	(2.2)	(10.3)	(23.3)	(30.0)	(21.1)	(9.6)	(2.6)	(0.8)	(100.0)				
9～	5	343	1,784	5,096	8,584	7,429	3,748	1,203	370	28,562	46	28,608	9.68	2.67
(%)	(0.0)	(1.2)	(6.2)	(17.8)	(30.1)	(26.0)	(13.1)	(4.2)	(1.3)	(100.0)				
10～	17	454	2,294	8,120	16,559	17,169	10,021	3,903	1,162	59,699	70	59,769	10.28	2.66
(%)	(0.0)	(0.8)	(3.8)	(13.6)	(27.7)	(28.8)	(16.8)	(6.5)	(1.9)	(100.0)				
11～	28	318	1,552	5,394	12,019	14,131	9,491	4,001	1,417	48,351	64	48,415	10.67	2.75
(%)	(0.1)	(0.7)	(3.2)	(11.2)	(24.9)	(29.2)	(19.6)	(8.3)	(2.9)	(100.0)				
12～	9	128	538	1,796	3,824	4,771	3,734	1,725	660	17,185	23	17,208	10.92	2.85
(%)	(0.1)	(0.7)	(3.1)	(10.5)	(22.3)	(27.8)	(21.7)	(10.0)	(3.8)	(100.0)				
13～	5	44	128	347	822	1,177	985	529	214	4,251	9	4,260	11.26	3.00
(%)	(0.1)	(1.0)	(3.0)	(8.2)	(19.3)	(27.7)	(23.2)	(12.4)	(5.0)	(100.0)				
14～	1	13	34	115	271	388	353	245	130	1,550	1	1,551	11.75	3.12
(%)	(0.1)	(0.8)	(2.2)	(7.4)	(17.5)	(25.0)	(22.8)	(15.8)	(8.4)	(100.0)				
合計	74	1,632	7,700	23,744	45,625	47,478	29,401	11,889	4,050	171,593	238	171,831	10.29	2.79
(%)	(0.0)	(1.0)	(4.5)	(13.8)	(26.6)	(27.7)	(17.1)	(6.9)	(2.4)	(100.0)				
記載なし	2	24	95	262	490	556	353	147	50	1,979	4,399	6,378	10.37	2.92
(%)	(0.1)	(1.2)	(4.8)	(13.2)	(24.8)	(28.1)	(17.8)	(7.4)	(2.5)	(100.0)				
総計	76	1,656	7,795	24,006	46,115	48,034	29,754	12,036	4,100	173,572	4,637	178,209	10.29	2.79
(%)	(0.0)	(1.0)	(4.5)	(13.8)	(26.6)	(27.7)	(17.1)	(6.9)	(2.4)	(100.0)				
平均	10.94	10.16	10.19	10.38	10.58	10.78	10.97	11.14	11.28	10.71	10.62	10.71		
標準偏差	1.46	1.60	1.43	1.31	1.22	1.18	1.18	1.19	1.29	1.25	1.34	1.25		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査