

表 2988 クレアチニン濃度(mg/dL) 体重(kg)別 (腹膜透析患者)

体重	2未満	2～	4～	6～	8～	10～	12～	14～	16～	合計	記載なし	総計	平均	標準偏差
20未満	1	6	15	9	5			1	1	38		38	6.08	3.14
(%)	(2.6)	(15.8)	(39.5)	(23.7)	(13.2)			(2.6)	(2.6)	(100.0)				
20～		2	5	1	4	6	3	2		23		23	8.97	3.60
(%)		(8.7)	(21.7)	(4.3)	(17.4)	(26.1)	(13.0)	(8.7)		(100.0)				
30～	1	15	40	57	33	13	6	1		166	6	172	7.13	2.57
(%)	(0.6)	(9.0)	(24.1)	(34.3)	(19.9)	(7.8)	(3.6)	(0.6)		(100.0)				
40～	2	43	127	212	246	140	70	21	13	874	30	904	8.49	2.90
(%)	(0.2)	(4.9)	(14.5)	(24.3)	(28.1)	(16.0)	(8.0)	(2.4)	(1.5)	(100.0)				
50～	4	34	156	340	386	306	216	69	37	1,548	43	1,591	9.39	3.17
(%)	(0.3)	(2.2)	(10.1)	(22.0)	(24.9)	(19.8)	(14.0)	(4.5)	(2.4)	(100.0)				
60～	1	19	108	252	340	344	249	164	96	1,573	43	1,616	10.53	3.44
(%)	(0.1)	(1.2)	(6.9)	(16.0)	(21.6)	(21.9)	(15.8)	(10.4)	(6.1)	(100.0)				
70～	1	7	45	99	141	152	159	110	69	783	28	811	11.21	3.56
(%)	(0.1)	(0.9)	(5.7)	(12.6)	(18.0)	(19.4)	(20.3)	(14.0)	(8.8)	(100.0)				
80～		2	13	27	37	54	58	55	37	283	15	298	12.20	3.76
(%)		(0.7)	(4.6)	(9.5)	(13.1)	(19.1)	(20.5)	(19.4)	(13.1)	(100.0)				
90～		1	8	7	15	13	22	18	14	98	6	104	11.96	3.87
(%)		(1.0)	(8.2)	(7.1)	(15.3)	(13.3)	(22.4)	(18.4)	(14.3)	(100.0)				
100～		2		1		3	3	4	6	19	1	20	13.32	4.79
(%)		(10.5)		(5.3)		(15.8)	(15.8)	(21.1)	(31.6)	(100.0)				
110～				1	1	2	1	3	1	9	1	10	12.08	3.21
(%)				(11.1)	(11.1)	(22.2)	(11.1)	(33.3)	(11.1)	(100.0)				
120～			1	2	1		1	1	2	8	1	9	11.95	5.53
(%)			(12.5)	(25.0)	(12.5)		(12.5)	(12.5)	(25.0)	(100.0)				
合計	10	131	518	1,008	1,209	1,033	788	449	276	5,422	174	5,596	9.96	3.53
(%)	(0.2)	(2.4)	(9.6)	(18.6)	(22.3)	(19.1)	(14.5)	(8.3)	(5.1)	(100.0)				
記載なし	7	23	57	125	140	82	60	39	19	552	2,793	3,345	9.16	3.47
(%)	(1.3)	(4.2)	(10.3)	(22.6)	(25.4)	(14.9)	(10.9)	(7.1)	(3.4)	(100.0)				
総計	17	154	575	1,133	1,349	1,115	848	488	295	5,974	2,967	8,941	9.89	3.53
(%)	(0.3)	(2.6)	(9.6)	(19.0)	(22.6)	(18.7)	(14.2)	(8.2)	(4.9)	(100.0)				
平均	49.11	50.78	54.73	56.88	58.69	61.66	64.73	68.90	70.66	60.67	63.20	60.74		
標準偏差	16.60	16.62	15.48	13.03	12.13	11.87	12.69	13.13	14.15	13.84	15.27	13.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査