

表 3005 single pool Kt/V(Kt/Vsp) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	25	30	180	937	4,246	11,962	15,781	11,416	6,139	3,820	54,536	4,911	59,447	1.54	0.30
(%)	(0.0)	(0.1)	(0.3)	(1.7)	(7.8)	(21.9)	(28.9)	(20.9)	(11.3)	(7.0)	(100.0)				
慢性腎盂腎炎	1	2	5	39	147	320	448	365	214	159	1,700	117	1,817	1.57	0.33
(%)	(0.1)	(0.1)	(0.3)	(2.3)	(8.6)	(18.8)	(26.4)	(21.5)	(12.6)	(9.4)	(100.0)				
急速進行性糸球体腎炎	1		6	22	97	206	318	280	165	97	1,192	89	1,281	1.57	0.31
(%)	(0.1)		(0.5)	(1.8)	(8.1)	(17.3)	(26.7)	(23.5)	(13.8)	(8.1)	(100.0)				
妊娠腎/妊娠中毒症			2	4	26	74	239	281	177	119	922	77	999	1.69	0.28
(%)			(0.2)	(0.4)	(2.8)	(8.0)	(25.9)	(30.5)	(19.2)	(12.9)	(100.0)				
その他の分類不能の腎炎	1	1	4	11	52	163	189	141	72	70	704	59	763	1.55	0.32
(%)	(0.1)	(0.1)	(0.6)	(1.6)	(7.4)	(23.2)	(26.8)	(20.0)	(10.2)	(9.9)	(100.0)				
多発性嚢胞腎	2	1	22	162	637	1,638	1,883	1,179	497	257	6,278	506	6,784	1.48	0.28
(%)	(0.0)	(0.0)	(0.4)	(2.6)	(10.1)	(26.1)	(30.0)	(18.8)	(7.9)	(4.1)	(100.0)				
腎硬化症	5	19	74	356	1,514	3,518	3,893	2,441	1,152	638	13,610	1,067	14,677	1.48	0.29
(%)	(0.0)	(0.1)	(0.5)	(2.6)	(11.1)	(25.8)	(28.6)	(17.9)	(8.5)	(4.7)	(100.0)				
悪性高血圧	2	3	5	41	170	349	356	240	115	71	1,352	113	1,465	1.48	0.30
(%)	(0.1)	(0.2)	(0.4)	(3.0)	(12.6)	(25.8)	(26.3)	(17.8)	(8.5)	(5.3)	(100.0)				
糖尿病性腎症	39	84	383	2,405	9,231	18,294	16,933	8,970	3,460	1,710	61,509	5,341	66,850	1.42	0.28
(%)	(0.1)	(0.1)	(0.6)	(3.9)	(15.0)	(29.7)	(27.5)	(14.6)	(5.6)	(2.8)	(100.0)				
SLE腎炎	1		2	17	56	157	249	291	199	148	1,120	95	1,215	1.65	0.33
(%)	(0.1)		(0.2)	(1.5)	(5.0)	(14.0)	(22.2)	(26.0)	(17.8)	(13.2)	(100.0)				
アミロイド腎			1	4	6	32	57	40	32	28	200	30	230	1.64	0.34
(%)			(0.5)	(2.0)	(3.0)	(16.0)	(28.5)	(20.0)	(16.0)	(14.0)	(100.0)				
痛風腎				17	74	185	199	111	38	26	650	55	705	1.46	0.26
(%)				(2.6)	(11.4)	(28.5)	(30.6)	(17.1)	(5.8)	(4.0)	(100.0)				
先天性代謝異常による腎不全				5	14	25	34	32	21	10	141	13	154	1.57	0.34
(%)				(3.5)	(9.9)	(17.7)	(24.1)	(22.7)	(14.9)	(7.1)	(100.0)				
腎・尿路結核				1	13	18	31	33	12	17	125	10	135	1.59	0.31
(%)				(0.8)	(10.4)	(14.4)	(24.8)	(26.4)	(9.6)	(13.6)	(100.0)				
腎・尿路結石	1			5	26	87	110	55	35	14	333	21	354	1.51	0.29
(%)	(0.3)			(1.5)	(7.8)	(26.1)	(33.0)	(16.5)	(10.5)	(4.2)	(100.0)				
腎・尿路腫瘍	1	2	5	10	56	123	131	64	31	18	441	32	473	1.45	0.30
(%)	(0.2)	(0.5)	(1.1)	(2.3)	(12.7)	(27.9)	(29.7)	(14.5)	(7.0)	(4.1)	(100.0)				
閉塞性尿路障害		1	4	15	33	89	89	75	39	37	382	32	414	1.53	0.34
(%)		(0.3)	(1.0)	(3.9)	(8.6)	(23.3)	(23.3)	(19.6)	(10.2)	(9.7)	(100.0)				
骨髄腫			1	1	10	18	27	10	8	2	77	11	88	1.46	0.26
(%)			(1.3)	(1.3)	(13.0)	(23.4)	(35.1)	(13.0)	(10.4)	(2.6)	(100.0)				
腎形成不全				6	33	57	89	78	40	29	332	27	359	1.57	0.31
(%)				(1.8)	(9.9)	(17.2)	(26.8)	(23.5)	(12.0)	(8.7)	(100.0)				
不明	17	23	62	352	1,432	3,251	3,792	2,573	1,250	744	13,496	1,334	14,830	1.50	0.30
(%)	(0.1)	(0.2)	(0.5)	(2.6)	(10.6)	(24.1)	(28.1)	(19.1)	(9.3)	(5.5)	(100.0)				
再導入	1		3	17	80	224	285	235	122	106	1,073	109	1,182	1.57	0.31
(%)	(0.1)		(0.3)	(1.6)	(7.5)	(20.9)	(26.6)	(21.9)	(11.4)	(9.9)	(100.0)				
その他	3	1	15	77	375	861	1,003	725	373	246	3,679	308	3,987	1.52	0.31
(%)	(0.1)	(0.0)	(0.4)	(2.1)	(10.2)	(23.4)	(27.3)	(19.7)	(10.1)	(6.7)	(100.0)				
合計	100	167	774	4,504	18,328	41,651	46,136	29,635	14,191	8,366	163,852	14,357	178,209	1.48	0.30
(%)	(0.1)	(0.1)	(0.5)	(2.7)	(11.2)	(25.4)	(28.2)	(18.1)	(8.7)	(5.1)	(100.0)				
記載なし															
(%)															
総計	100	167	774	4,504	18,328	41,651	46,136	29,635	14,191	8,366	163,852	14,357	178,209	1.48	0.30
(%)	(0.1)	(0.1)	(0.5)	(2.7)	(11.2)	(25.4)	(28.2)	(18.1)	(8.7)	(5.1)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査