

表 3019 single pool Kt/V(Kt/Vsp) 透析前リン濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
3未満	9	20	89	298	762	1,462	1,576	1,103	570	380	6,269	620	6,889	1.47	0.33
(%)	(0.1)	(0.3)	(1.4)	(4.8)	(12.2)	(23.3)	(25.1)	(17.6)	(9.1)	(6.1)	(100.0)				
3.0～	10	16	48	247	833	1,826	2,080	1,444	678	475	7,657	579	8,236	1.49	0.31
(%)	(0.1)	(0.2)	(0.6)	(3.2)	(10.9)	(23.8)	(27.2)	(18.9)	(8.9)	(6.2)	(100.0)				
3.5～	11	17	60	363	1,363	3,181	3,704	2,498	1,268	836	13,301	893	14,194	1.50	0.31
(%)	(0.1)	(0.1)	(0.5)	(2.7)	(10.2)	(23.9)	(27.8)	(18.8)	(9.5)	(6.3)	(100.0)				
4.0～	10	18	100	504	2,083	4,820	5,753	3,805	1,919	1,189	20,201	1,251	21,452	1.50	0.30
(%)	(0.0)	(0.1)	(0.5)	(2.5)	(10.3)	(23.9)	(28.5)	(18.8)	(9.5)	(5.9)	(100.0)				
4.5～	14	18	94	551	2,430	5,890	7,183	4,749	2,320	1,424	24,673	1,405	26,078	1.51	0.29
(%)	(0.1)	(0.1)	(0.4)	(2.2)	(9.8)	(23.9)	(29.1)	(19.2)	(9.4)	(5.8)	(100.0)				
5.0～	12	15	95	593	2,606	6,410	7,260	4,728	2,232	1,255	25,206	1,418	26,624	1.49	0.29
(%)	(0.0)	(0.1)	(0.4)	(2.4)	(10.3)	(25.4)	(28.8)	(18.8)	(8.9)	(5.0)	(100.0)				
5.5～	12	18	77	547	2,300	5,736	6,290	4,113	1,869	1,056	22,018	1,104	23,122	1.49	0.29
(%)	(0.1)	(0.1)	(0.3)	(2.5)	(10.4)	(26.1)	(28.6)	(18.7)	(8.5)	(4.8)	(100.0)				
6.0～		1	9	80	427	994	1,076	666	326	161	3,740	226	3,966	1.48	0.28
(%)		(0.0)	(0.2)	(2.1)	(11.4)	(26.6)	(28.8)	(17.8)	(8.7)	(4.3)	(100.0)				
6.1～	6	11	47	327	1,492	3,335	3,642	2,093	1,006	551	12,510	655	13,165	1.47	0.29
(%)	(0.0)	(0.1)	(0.4)	(2.6)	(11.9)	(26.7)	(29.1)	(16.7)	(8.0)	(4.4)	(100.0)				
6.5～	3	8	53	297	1,360	3,035	2,979	1,793	829	413	10,770	592	11,362	1.46	0.28
(%)	(0.0)	(0.1)	(0.5)	(2.8)	(12.6)	(28.2)	(27.7)	(16.6)	(7.7)	(3.8)	(100.0)				
7.0～	3	3	27	200	905	1,905	1,854	1,042	477	271	6,687	393	7,080	1.45	0.29
(%)	(0.0)	(0.0)	(0.4)	(3.0)	(13.5)	(28.5)	(27.7)	(15.6)	(7.1)	(4.1)	(100.0)				
7.5～	3	11	22	158	572	1,135	1,073	600	266	129	3,969	229	4,198	1.43	0.28
(%)	(0.1)	(0.3)	(0.6)	(4.0)	(14.4)	(28.6)	(27.0)	(15.1)	(6.7)	(3.3)	(100.0)				
8.0～	2	5	16	97	401	666	600	337	159	75	2,358	127	2,485	1.41	0.29
(%)	(0.1)	(0.2)	(0.7)	(4.1)	(17.0)	(28.2)	(25.4)	(14.3)	(6.7)	(3.2)	(100.0)				
8.5～	1	3	11	80	277	434	364	215	97	46	1,528	73	1,601	1.40	0.29
(%)	(0.1)	(0.2)	(0.7)	(5.2)	(18.1)	(28.4)	(23.8)	(14.1)	(6.3)	(3.0)	(100.0)				
9.0～	1	2	22	142	446	675	553	338	134	81	2,394	143	2,537	1.40	0.30
(%)	(0.0)	(0.1)	(0.9)	(5.9)	(18.6)	(28.2)	(23.1)	(14.1)	(5.6)	(3.4)	(100.0)				
合計	97	166	770	4,484	18,257	41,504	45,987	29,524	14,150	8,342	163,281	9,708	172,989	1.48	0.30
(%)	(0.1)	(0.1)	(0.5)	(2.7)	(11.2)	(25.4)	(28.2)	(18.1)	(8.7)	(5.1)	(100.0)				
記載なし	3	1	4	20	71	147	149	111	41	24	571	4,649	5,220	1.45	0.31
(%)	(0.5)	(0.2)	(0.7)	(3.5)	(12.4)	(25.7)	(26.1)	(19.4)	(7.2)	(4.2)	(100.0)				
総計	100	167	774	4,504	18,328	41,651	46,136	29,635	14,191	8,366	163,852	14,357	178,209	1.48	0.30
(%)	(0.1)	(0.1)	(0.5)	(2.7)	(11.2)	(25.4)	(28.2)	(18.1)	(8.7)	(5.1)	(100.0)				
平均	4.84	4.96	5.05	5.30	5.39	5.32	5.22	5.15	5.10	5.00	5.23	5.07	5.22		
標準偏差	1.59	1.79	1.81	1.72	1.58	1.44	1.37	1.35	1.34	1.33	1.42	1.50	1.43		

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査