

表 3059 equilibrated Kt/V(eKt/V) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	33	79	605	3,310	11,981	17,421	12,230	5,728	1,983	1,188	54,558	4,889	59,447	1.35	0.28
(%)	(0.1)	(0.1)	(1.1)	(6.1)	(22.0)	(31.9)	(22.4)	(10.5)	(3.6)	(2.2)	(100.0)				
慢性腎盂腎炎	1	4	28	116	332	501	382	199	80	58	1,701	116	1,817	1.37	0.31
(%)	(0.1)	(0.2)	(1.6)	(6.8)	(19.5)	(29.5)	(22.5)	(11.7)	(4.7)	(3.4)	(100.0)				
急速進行性糸球体腎炎	1	2	18	80	228	324	311	152	56	19	1,191	90	1,281	1.37	0.28
(%)	(0.1)	(0.2)	(1.5)	(6.7)	(19.1)	(27.2)	(26.1)	(12.8)	(4.7)	(1.6)	(100.0)				
妊娠腎/妊娠中毒症			4	28	76	251	310	169	51	34	923	76	999	1.47	0.26
(%)			(0.4)	(3.0)	(8.2)	(27.2)	(33.6)	(18.3)	(5.5)	(3.7)	(100.0)				
その他の分類不能の腎炎	2	1	10	43	162	202	155	74	28	27	704	59	763	1.36	0.30
(%)	(0.3)	(0.1)	(1.4)	(6.1)	(23.0)	(28.7)	(22.0)	(10.5)	(4.0)	(3.8)	(100.0)				
多発性嚢胞腎	2	4	105	525	1,637	2,095	1,235	448	159	70	6,280	504	6,784	1.29	0.25
(%)	(0.0)	(0.1)	(1.7)	(8.4)	(26.1)	(33.4)	(19.7)	(7.1)	(2.5)	(1.1)	(100.0)				
腎硬化症	8	40	247	1,323	3,533	4,348	2,537	1,076	333	172	13,617	1,060	14,677	1.29	0.27
(%)	(0.1)	(0.3)	(1.8)	(9.7)	(25.9)	(31.9)	(18.6)	(7.9)	(2.4)	(1.3)	(100.0)				
悪性高血圧	3	2	28	117	372	408	255	112	35	21	1,353	112	1,465	1.29	0.28
(%)	(0.2)	(0.1)	(2.1)	(8.6)	(27.5)	(30.2)	(18.8)	(8.3)	(2.6)	(1.6)	(100.0)				
糖尿病性腎症	56	174	1,489	7,570	18,901	19,355	9,408	3,164	919	497	61,533	5,317	66,850	1.23	0.25
(%)	(0.1)	(0.3)	(2.4)	(12.3)	(30.7)	(31.5)	(15.3)	(5.1)	(1.5)	(0.8)	(100.0)				
SLE腎炎	1		10	51	166	282	301	186	77	47	1,121	94	1,215	1.44	0.31
(%)	(0.1)		(0.9)	(4.5)	(14.8)	(25.2)	(26.9)	(16.6)	(6.9)	(4.2)	(100.0)				
アミロイド腎		1	1	7	37	57	46	30	12	9	200	30	230	1.42	0.31
(%)		(0.5)	(0.5)	(3.5)	(18.5)	(28.5)	(23.0)	(15.0)	(6.0)	(4.5)	(100.0)				
痛風腎			4	56	191	231	110	36	15	7	650	55	705	1.27	0.24
(%)			(0.6)	(8.6)	(29.4)	(35.5)	(16.9)	(5.5)	(2.3)	(1.1)	(100.0)				
先天性代謝異常による腎不全			2	14	27	35	36	17	4	6	141	13	154	1.37	0.31
(%)			(1.4)	(9.9)	(19.1)	(24.8)	(25.5)	(12.1)	(2.8)	(4.3)	(100.0)				
腎・尿路結核			1	9	20	34	36	15	8	2	125	10	135	1.39	0.29
(%)			(0.8)	(7.2)	(16.0)	(27.2)	(28.8)	(12.0)	(6.4)	(1.6)	(100.0)				
腎・尿路結石	1		3	23	78	129	56	27	10	6	333	21	354	1.31	0.26
(%)	(0.3)		(0.9)	(6.9)	(23.4)	(38.7)	(16.8)	(8.1)	(3.0)	(1.8)	(100.0)				
腎・尿路腫瘍	1	4	5	48	124	151	64	30	9	5	441	32	473	1.26	0.27
(%)	(0.2)	(0.9)	(1.1)	(10.9)	(28.1)	(34.2)	(14.5)	(6.8)	(2.0)	(1.1)	(100.0)				
閉塞性尿路障害		3	11	28	83	111	75	43	16	12	382	32	414	1.33	0.31
(%)		(0.8)	(2.9)	(7.3)	(21.7)	(29.1)	(19.6)	(11.3)	(4.2)	(3.1)	(100.0)				
骨髄腫			2	8	18	30	10	8	1		77	11	88	1.26	0.24
(%)			(2.6)	(10.4)	(23.4)	(39.0)	(13.0)	(10.4)	(1.3)		(100.0)				
腎形成不全			4	27	56	94	87	45	10	10	333	26	359	1.38	0.29
(%)			(1.2)	(8.1)	(16.8)	(28.2)	(26.1)	(13.5)	(3.0)	(3.0)	(100.0)				
不明	21	43	204	1,276	3,280	4,258	2,676	1,176	375	196	13,505	1,325	14,830	1.30	0.28
(%)	(0.2)	(0.3)	(1.5)	(9.4)	(24.3)	(31.5)	(19.8)	(8.7)	(2.8)	(1.5)	(100.0)				
再導入	1	1	12	62	231	309	248	115	64	31	1,074	108	1,182	1.37	0.29
(%)	(0.1)	(0.1)	(1.1)	(5.8)	(21.5)	(28.8)	(23.1)	(10.7)	(6.0)	(2.9)	(100.0)				
その他	3	8	45	319	862	1,142	755	347	132	71	3,684	303	3,987	1.32	0.28
(%)	(0.1)	(0.2)	(1.2)	(8.7)	(23.4)	(31.0)	(20.5)	(9.4)	(3.6)	(1.9)	(100.0)				
合計	134	366	2,838	15,040	42,395	51,768	31,323	13,197	4,377	2,488	163,926	14,283	178,209	1.29	0.27
(%)	(0.1)	(0.2)	(1.7)	(9.2)	(25.9)	(31.6)	(19.1)	(8.1)	(2.7)	(1.5)	(100.0)				
記載なし															
(%)															
総計	134	366	2,838	15,040	42,395	51,768	31,323	13,197	4,377	2,488	163,926	14,283	178,209	1.29	0.27
(%)	(0.1)	(0.2)	(1.7)	(9.2)	(25.9)	(31.6)	(19.1)	(8.1)	(2.7)	(1.5)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査