

表 3073 equilibrated Kt/V(eKt/V) 透析前リン濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
3未満	11	48	215	749	1,479	1,739	1,169	525	203	135	6,273	616	6,889	1.28	0.31
(%)	(0.2)	(0.8)	(3.4)	(11.9)	(23.6)	(27.7)	(18.6)	(8.4)	(3.2)	(2.2)	(100.0)				
3.0～	15	30	159	715	1,853	2,295	1,557	654	232	155	7,665	571	8,236	1.30	0.29
(%)	(0.2)	(0.4)	(2.1)	(9.3)	(24.2)	(29.9)	(20.3)	(8.5)	(3.0)	(2.0)	(100.0)				
3.5～	16	29	242	1,142	3,217	4,096	2,664	1,202	431	268	13,307	887	14,194	1.31	0.28
(%)	(0.1)	(0.2)	(1.8)	(8.6)	(24.2)	(30.8)	(20.0)	(9.0)	(3.2)	(2.0)	(100.0)				
4.0～	14	49	359	1,729	4,842	6,422	3,983	1,808	617	388	20,211	1,241	21,452	1.31	0.28
(%)	(0.1)	(0.2)	(1.8)	(8.6)	(24.0)	(31.8)	(19.7)	(8.9)	(3.1)	(1.9)	(100.0)				
4.5～	18	43	347	1,980	5,862	8,051	5,002	2,212	766	402	24,683	1,395	26,078	1.31	0.27
(%)	(0.1)	(0.2)	(1.4)	(8.0)	(23.7)	(32.6)	(20.3)	(9.0)	(3.1)	(1.6)	(100.0)				
5.0～	15	39	387	2,072	6,508	8,124	4,977	2,040	685	373	25,220	1,404	26,624	1.30	0.27
(%)	(0.1)	(0.2)	(1.5)	(8.2)	(25.8)	(32.2)	(19.7)	(8.1)	(2.7)	(1.5)	(100.0)				
5.5～	13	40	320	1,870	5,768	7,092	4,367	1,713	532	309	22,024	1,098	23,122	1.30	0.26
(%)	(0.1)	(0.2)	(1.5)	(8.5)	(26.2)	(32.2)	(19.8)	(7.8)	(2.4)	(1.4)	(100.0)				
6.0～		5	45	315	1,045	1,216	702	291	86	37	3,742	224	3,966	1.29	0.25
(%)		(0.1)	(1.2)	(8.4)	(27.9)	(32.5)	(18.8)	(7.8)	(2.3)	(1.0)	(100.0)				
6.1～	6	20	203	1,179	3,429	4,092	2,233	888	308	154	12,512	653	13,165	1.28	0.26
(%)	(0.0)	(0.2)	(1.6)	(9.4)	(27.4)	(32.7)	(17.8)	(7.1)	(2.5)	(1.2)	(100.0)				
6.5～	6	16	179	1,082	3,147	3,359	1,909	770	201	107	10,776	586	11,362	1.27	0.26
(%)	(0.1)	(0.1)	(1.7)	(10.0)	(29.2)	(31.2)	(17.7)	(7.1)	(1.9)	(1.0)	(100.0)				
7.0～	3	8	106	732	1,979	2,109	1,099	446	134	74	6,690	390	7,080	1.26	0.26
(%)	(0.0)	(0.1)	(1.6)	(10.9)	(29.6)	(31.5)	(16.4)	(6.7)	(2.0)	(1.1)	(100.0)				
7.5～	6	13	82	466	1,185	1,240	646	234	65	32	3,969	229	4,198	1.24	0.26
(%)	(0.2)	(0.3)	(2.1)	(11.7)	(29.9)	(31.2)	(16.3)	(5.9)	(1.6)	(0.8)	(100.0)				
8.0～	3	10	58	330	711	692	351	146	44	13	2,358	127	2,485	1.23	0.26
(%)	(0.1)	(0.4)	(2.5)	(14.0)	(30.2)	(29.3)	(14.9)	(6.2)	(1.9)	(0.6)	(100.0)				
8.5～	3	4	47	239	462	426	222	90	27	9	1,529	72	1,601	1.22	0.26
(%)	(0.2)	(0.3)	(3.1)	(15.6)	(30.2)	(27.9)	(14.5)	(5.9)	(1.8)	(0.6)	(100.0)				
9.0～	1	9	79	375	755	642	331	139	39	26	2,396	141	2,537	1.22	0.27
(%)	(0.0)	(0.4)	(3.3)	(15.7)	(31.5)	(26.8)	(13.8)	(5.8)	(1.6)	(1.1)	(100.0)				
合計	130	363	2,828	14,975	42,242	51,595	31,212	13,158	4,370	2,482	163,355	9,634	172,989	1.29	0.27
(%)	(0.1)	(0.2)	(1.7)	(9.2)	(25.9)	(31.6)	(19.1)	(8.1)	(2.7)	(1.5)	(100.0)				
記載なし	4	3	10	65	153	173	111	39	7	6	571	4,649	5,220	1.25	0.27
(%)	(0.7)	(0.5)	(1.8)	(11.4)	(26.8)	(30.3)	(19.4)	(6.8)	(1.2)	(1.1)	(100.0)				
総計	134	366	2,838	15,040	42,395	51,768	31,323	13,197	4,377	2,488	163,926	14,283	178,209	1.29	0.27
(%)	(0.1)	(0.2)	(1.7)	(9.2)	(25.9)	(31.6)	(19.1)	(8.1)	(2.7)	(1.5)	(100.0)				
平均	4.87	4.92	5.18	5.35	5.34	5.23	5.15	5.09	5.00	4.90	5.23	5.07	5.22		
標準偏差	1.69	1.74	1.71	1.62	1.46	1.37	1.34	1.34	1.32	1.33	1.42	1.50	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査