

表 3113 nPCR(g/kg/day) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	85	1,997	14,044	24,555	11,492	2,156	237	32	11	14	54,623	4,824	59,447	0.89	0.18
(%)	(0.2)	(3.7)	(25.7)	(45.0)	(21.0)	(3.9)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
慢性腎盂腎炎	5	83	427	759	353	71	6	1			1,705	112	1,817	0.88	0.18
(%)	(0.3)	(4.9)	(25.0)	(44.5)	(20.7)	(4.2)	(0.4)	(0.1)			(100.0)				
急速進行性糸球体腎炎	5	65	314	503	254	46	6				1,193	88	1,281	0.88	0.18
(%)	(0.4)	(5.4)	(26.3)	(42.2)	(21.3)	(3.9)	(0.5)				(100.0)				
妊娠腎/妊娠中毒症	1	25	197	412	228	51	9	2			925	74	999	0.92	0.18
(%)	(0.1)	(2.7)	(21.3)	(44.5)	(24.6)	(5.5)	(1.0)	(0.2)			(100.0)				
その他の分類不能の腎炎	3	21	186	315	147	26	6				704	59	763	0.89	0.18
(%)	(0.4)	(3.0)	(26.4)	(44.7)	(20.9)	(3.7)	(0.9)				(100.0)				
多発性嚢胞腎	10	196	1,760	2,983	1,163	156	11	2	1	3	6,285	499	6,784	0.87	0.16
(%)	(0.2)	(3.1)	(28.0)	(47.5)	(18.5)	(2.5)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
腎硬化症	31	730	4,444	5,762	2,261	363	35	6	2		13,634	1,043	14,677	0.85	0.17
(%)	(0.2)	(5.4)	(32.6)	(42.3)	(16.6)	(2.7)	(0.3)	(0.0)	(0.0)		(100.0)				
悪性高血圧	2	68	382	580	257	57	7	1	1	1	1,356	109	1,465	0.88	0.19
(%)	(0.1)	(5.0)	(28.2)	(42.8)	(19.0)	(4.2)	(0.5)	(0.1)	(0.1)	(0.1)	(100.0)				
糖尿病性腎症	168	4,122	22,846	25,205	7,899	1,183	128	23	5	16	61,595	5,255	66,850	0.83	0.17
(%)	(0.3)	(6.7)	(37.1)	(40.9)	(12.8)	(1.9)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
SLE腎炎	1	39	241	483	282	66	15				1,127	88	1,215	0.92	0.19
(%)	(0.1)	(3.5)	(21.4)	(42.9)	(25.0)	(5.9)	(1.3)				(100.0)				
アミロイド腎	1	11	55	94	34	5					200	30	230	0.85	0.17
(%)	(0.5)	(5.5)	(27.5)	(47.0)	(17.0)	(2.5)					(100.0)				
痛風腎	1	20	180	270	154	21	4			1	651	54	705	0.89	0.19
(%)	(0.2)	(3.1)	(27.6)	(41.5)	(23.7)	(3.2)	(0.6)			(0.2)	(100.0)				
先天性代謝異常による腎不全		7	38	51	37	6	2				141	13	154	0.91	0.19
(%)		(5.0)	(27.0)	(36.2)	(26.2)	(4.3)	(1.4)				(100.0)				
腎・尿路結核		5	38	54	23	5					125	10	135	0.88	0.17
(%)		(4.0)	(30.4)	(43.2)	(18.4)	(4.0)					(100.0)				
腎・尿路結石	2	12	95	157	52	15					333	21	354	0.87	0.17
(%)	(0.6)	(3.6)	(28.5)	(47.1)	(15.6)	(4.5)					(100.0)				
腎・尿路腫瘍	1	21	146	190	75	7	1				441	32	473	0.85	0.16
(%)	(0.2)	(4.8)	(33.1)	(43.1)	(17.0)	(1.6)	(0.2)				(100.0)				
閉塞性尿路障害	3	23	111	152	74	17	2	1			383	31	414	0.87	0.20
(%)	(0.8)	(6.0)	(29.0)	(39.7)	(19.3)	(4.4)	(0.5)	(0.3)			(100.0)				
骨髄腫		8	21	33	14	1					77	11	88	0.84	0.17
(%)		(10.4)	(27.3)	(42.9)	(18.2)	(1.3)					(100.0)				
腎形成不全	1	3	61	166	82	17	3				333	26	359	0.93	0.16
(%)	(0.3)	(0.9)	(18.3)	(49.8)	(24.6)	(5.1)	(0.9)				(100.0)				
不明	34	746	4,072	5,760	2,367	465	61	6	2	1	13,514	1,316	14,830	0.86	0.18
(%)	(0.3)	(5.5)	(30.1)	(42.6)	(17.5)	(3.4)	(0.5)	(0.0)	(0.0)	(0.0)	(100.0)				
再導入	1	18	214	500	282	56	6		1		1,078	104	1,182	0.92	0.17
(%)	(0.1)	(1.7)	(19.9)	(46.4)	(26.2)	(5.2)	(0.6)		(0.1)		(100.0)				
その他	14	198	1,084	1,527	718	126	16	5	2	2	3,692	295	3,987	0.87	0.19
(%)	(0.4)	(5.4)	(29.4)	(41.4)	(19.4)	(3.4)	(0.4)	(0.1)	(0.1)	(0.1)	(100.0)				
合計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし															
(%)															
総計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査