

表 3114 nPCR(g/kg/day) 医療機関別（施設血液透析患者 透析回数週3回 透析歴2年以上）

医療機関	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	3	32	56	54	15	3					163	77	240	0.77	0.19
(%)	(1.8)	(19.6)	(34.4)	(33.1)	(9.2)	(1.8)					(100.0)				
私立大学	6	36	288	360	136	27	2				855	164	1,019	0.85	0.18
(%)	(0.7)	(4.2)	(33.7)	(42.1)	(15.9)	(3.2)	(0.2)				(100.0)				
国立	2	26	107	119	35	6	2				297	34	331	0.82	0.18
(%)	(0.7)	(8.8)	(36.0)	(40.1)	(11.8)	(2.0)	(0.7)				(100.0)				
県市町村立	19	571	2,964	3,617	1,354	218	29	5		2	8,779	967	9,746	0.84	0.18
(%)	(0.2)	(6.5)	(33.8)	(41.2)	(15.4)	(2.5)	(0.3)	(0.1)		(0.0)	(100.0)				
社会保険	6	103	628	857	319	55	7	4	1	1	1,981	172	2,153	0.86	0.18
(%)	(0.3)	(5.2)	(31.7)	(43.3)	(16.1)	(2.8)	(0.4)	(0.2)	(0.1)	(0.1)	(100.0)				
厚生連	14	288	1,631	1,972	729	107	14	2		2	4,759	337	5,096	0.84	0.17
(%)	(0.3)	(6.1)	(34.3)	(41.4)	(15.3)	(2.2)	(0.3)	(0.0)		(0.0)	(100.0)				
その他公的	11	294	1,782	2,144	771	126	20	1		1	5,150	433	5,583	0.84	0.17
(%)	(0.2)	(5.7)	(34.6)	(41.6)	(15.0)	(2.4)	(0.4)	(0.0)		(0.0)	(100.0)				
私立総合	7	211	1,094	1,384	586	86	12	2	3	1	3,386	227	3,613	0.85	0.18
(%)	(0.2)	(6.2)	(32.3)	(40.9)	(17.3)	(2.5)	(0.4)	(0.1)	(0.1)	(0.0)	(100.0)				
私立	175	3,190	15,420	19,219	7,211	1,312	151	27	7	10	46,722	4,892	51,614	0.84	0.18
(%)	(0.4)	(6.8)	(33.0)	(41.1)	(15.4)	(2.8)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
私立診療所	126	3,667	26,986	40,785	17,092	2,976	318	38	14	21	92,023	6,791	98,814	0.87	0.17
(%)	(0.1)	(4.0)	(29.3)	(44.3)	(18.6)	(3.2)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
総計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査