

表 3119 nPCR(g/kg/day) %クレアチニン産生速度(%)別（施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上）

%クレアチニン 産生速度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
60未満	129	1,629	3,461	1,699	397	92	10	5	1	1	7,424		7,424	0.72	0.18
(%)	(1.7)	(21.9)	(46.6)	(22.9)	(5.3)	(1.2)	(0.1)	(0.1)	(0.0)	(0.0)	(100.0)				
60～	19	789	2,720	1,765	428	56	9				5,786		5,786	0.76	0.16
(%)	(0.3)	(13.6)	(47.0)	(30.5)	(7.4)	(1.0)	(0.2)				(100.0)				
70～	21	847	3,949	3,142	833	103	13	1			8,909		8,909	0.79	0.16
(%)	(0.2)	(9.5)	(44.3)	(35.3)	(9.4)	(1.2)	(0.1)	(0.0)			(100.0)				
80～	11	764	5,018	5,124	1,400	220	20	2			12,559		12,559	0.82	0.16
(%)	(0.1)	(6.1)	(40.0)	(40.8)	(11.1)	(1.8)	(0.2)	(0.0)			(100.0)				
90～	14	612	5,726	7,113	2,172	314	36	1	1		15,989		15,989	0.84	0.16
(%)	(0.1)	(3.8)	(35.8)	(44.5)	(13.6)	(2.0)	(0.2)	(0.0)	(0.0)		(100.0)				
100～	6	395	5,135	7,964	2,856	420	38	3		1	16,818		16,818	0.87	0.15
(%)	(0.0)	(2.3)	(30.5)	(47.4)	(17.0)	(2.5)	(0.2)	(0.0)		(0.0)	(100.0)				
110～	15	243	3,858	7,521	2,939	434	49	4			15,063		15,063	0.89	0.15
(%)	(0.1)	(1.6)	(25.6)	(49.9)	(19.5)	(2.9)	(0.3)	(0.0)			(100.0)				
120～	8	111	2,400	5,494	2,445	330	23	8			10,819		10,819	0.91	0.15
(%)	(0.1)	(1.0)	(22.2)	(50.8)	(22.6)	(3.1)	(0.2)	(0.1)			(100.0)				
130～	5	38	1,166	3,063	1,540	258	24	3	1	1	6,099		6,099	0.92	0.15
(%)	(0.1)	(0.6)	(19.1)	(50.2)	(25.3)	(4.2)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
140～	8	25	690	2,081	1,086	201	43	6	7	3	4,150	1	4,151	0.94	0.17
(%)	(0.2)	(0.6)	(16.6)	(50.1)	(26.2)	(4.8)	(1.0)	(0.1)	(0.2)	(0.1)	(100.0)				
合計	236	5,453	34,123	44,966	16,096	2,428	265	33	10	6	103,616	1	103,617	0.85	0.17
(%)	(0.2)	(5.3)	(32.9)	(43.4)	(15.5)	(2.3)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	2	2	10	9	6		7	1	4	22	63	8,835	8,898	1.52	0.75
(%)	(3.2)	(3.2)	(15.9)	(14.3)	(9.5)		(11.1)	(1.6)	(6.3)	(34.9)	(100.0)				
総計	238	5,455	34,133	44,975	16,102	2,428	272	34	14	28	103,679	8,836	112,515	0.85	0.17
(%)	(0.2)	(5.3)	(32.9)	(43.4)	(15.5)	(2.3)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
平均	64.46	72.88	92.12	103.46	108.58	109.12	113.04	115.24	183.72	167.65	99.00	225.10	99.00		
標準偏差	35.70	26.34	24.93	23.13	22.76	25.58	32.66	46.58	77.73	84.91	25.64		25.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査