

表 3122 nPCR(g/kg/day) 透析前CRP濃度(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前CRP濃度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
0.2未満	82	2,699	22,586	35,040	14,711	2,543	253	35	10	16	77,975	3,531	81,506	0.88	0.17
(%)	(0.1)	(3.5)	(29.0)	(44.9)	(18.9)	(3.3)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
0.2～	66	1,502	8,559	11,412	4,131	687	72	15	4	6	26,454	1,386	27,840	0.85	0.17
(%)	(0.2)	(5.7)	(32.4)	(43.1)	(15.6)	(2.6)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
0.5～	46	962	4,709	5,404	1,944	293	42	4	2	5	13,411	835	14,246	0.84	0.18
(%)	(0.3)	(7.2)	(35.1)	(40.3)	(14.5)	(2.2)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
1.0～	39	866	3,103	3,234	1,158	190	29	2	2	2	8,625	620	9,245	0.82	0.18
(%)	(0.5)	(10.0)	(36.0)	(37.5)	(13.4)	(2.2)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
2.0～	39	561	1,904	1,843	643	129	18	3	2		5,142	501	5,643	0.81	0.19
(%)	(0.8)	(10.9)	(37.0)	(35.8)	(12.5)	(2.5)	(0.4)	(0.1)	(0.0)		(100.0)				
4.0～	18	206	668	612	218	36	5	1			1,764	212	1,976	0.80	0.19
(%)	(1.0)	(11.7)	(37.9)	(34.7)	(12.4)	(2.0)	(0.3)	(0.1)			(100.0)				
6.0～	4	103	298	288	99	26	1			1	820	129	949	0.81	0.20
(%)	(0.5)	(12.6)	(36.3)	(35.1)	(12.1)	(3.2)	(0.1)			(0.1)	(100.0)				
8.0～	6	63	165	152	54	13	3				456	76	532	0.80	0.20
(%)	(1.3)	(13.8)	(36.2)	(33.3)	(11.8)	(2.9)	(0.7)				(100.0)				
10.0～	5	50	222	186	76	13	4	2			558	93	651	0.82	0.20
(%)	(0.9)	(9.0)	(39.8)	(33.3)	(13.6)	(2.3)	(0.7)	(0.4)			(100.0)				
15.0～	5	40	127	125	44	17	1	2			361	88	449	0.82	0.21
(%)	(1.4)	(11.1)	(35.2)	(34.6)	(12.2)	(4.7)	(0.3)	(0.6)			(100.0)				
合計	310	7,052	42,341	58,296	23,078	3,947	428	64	20	30	135,566	7,471	143,037	0.86	0.17
(%)	(0.2)	(5.2)	(31.2)	(43.0)	(17.0)	(2.9)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	59	1,366	8,615	12,215	5,170	969	127	15	5	8	28,549	6,623	35,172	0.87	0.18
(%)	(0.2)	(4.8)	(30.2)	(42.8)	(18.1)	(3.4)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
総計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
平均	1.85	1.12	0.70	0.53	0.48	0.57	0.68	1.35	0.57	0.56	0.61	1.26	0.64		
標準偏差	3.67	2.55	2.04	1.64	1.60	2.18	1.87	3.77	0.80	1.20	1.86	3.35	1.97		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査