

表 3124 nPCR(g/kg/day) 透析前リン濃度(mg/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
3未満	146	1,673	2,637	1,399	346	65	7	5	2	3	6,283	606	6,889	0.71	0.19
(%)	(2.3)	(26.6)	(42.0)	(22.3)	(5.5)	(1.0)	(0.1)	(0.1)	(0.0)	(0.0)	(100.0)				
3.0～	52	1,368	3,405	2,204	562	82	7	1			7,681	555	8,236	0.75	0.17
(%)	(0.7)	(17.8)	(44.3)	(28.7)	(7.3)	(1.1)	(0.1)	(0.0)			(100.0)				
3.5～	27	1,487	5,934	4,541	1,180	140	16	2	1	5	13,333	861	14,194	0.78	0.17
(%)	(0.2)	(11.2)	(44.5)	(34.1)	(8.9)	(1.1)	(0.1)	(0.0)	(0.0)	(0.0)	(100.0)				
4.0～	35	1,434	8,329	7,883	2,247	266	30	2	3	2	20,231	1,221	21,452	0.81	0.16
(%)	(0.2)	(7.1)	(41.2)	(39.0)	(11.1)	(1.3)	(0.1)	(0.0)	(0.0)	(0.0)	(100.0)				
4.5～	29	949	8,914	10,949	3,395	422	41	4	1	2	24,706	1,372	26,078	0.84	0.16
(%)	(0.1)	(3.8)	(36.1)	(44.3)	(13.7)	(1.7)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
5.0～	27	654	7,892	11,922	4,028	659	47	8	2	8	25,247	1,377	26,624	0.87	0.16
(%)	(0.1)	(2.6)	(31.3)	(47.2)	(16.0)	(2.6)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
5.5～	18	400	5,778	10,717	4,384	685	51	9	3	5	22,050	1,072	23,122	0.89	0.16
(%)	(0.1)	(1.8)	(26.2)	(48.6)	(19.9)	(3.1)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
6.0～		60	897	1,821	817	136	14	2			3,747	219	3,966	0.90	0.16
(%)		(1.6)	(23.9)	(48.6)	(21.8)	(3.6)	(0.4)	(0.1)			(100.0)				
6.1～	7	133	2,789	6,139	2,903	498	49	4	2	3	12,527	638	13,165	0.91	0.16
(%)	(0.1)	(1.1)	(22.3)	(49.0)	(23.2)	(4.0)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
6.5～	8	95	1,960	5,281	2,860	524	47	5	2	2	10,784	578	11,362	0.93	0.16
(%)	(0.1)	(0.9)	(18.2)	(49.0)	(26.5)	(4.9)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
7.0～	6	49	1,042	3,168	1,972	390	55	7	3	2	6,694	386	7,080	0.95	0.17
(%)	(0.1)	(0.7)	(15.6)	(47.3)	(29.5)	(5.8)	(0.8)	(0.1)	(0.0)	(0.0)	(100.0)				
7.5～	3	30	524	1,787	1,267	316	39	5		2	3,973	225	4,198	0.97	0.17
(%)	(0.1)	(0.8)	(13.2)	(45.0)	(31.9)	(8.0)	(1.0)	(0.1)		(0.1)	(100.0)				
8.0～	3	18	272	990	801	228	39	7	1	1	2,360	125	2,485	0.99	0.18
(%)	(0.1)	(0.8)	(11.5)	(41.9)	(33.9)	(9.7)	(1.7)	(0.3)	(0.0)	(0.0)	(100.0)				
8.5～		15	198	602	534	153	26	3			1,531	70	1,601	0.99	0.18
(%)		(1.0)	(12.9)	(39.3)	(34.9)	(10.0)	(1.7)	(0.2)			(100.0)				
9.0～	2	27	226	860	838	337	84	15	5	3	2,397	140	2,537	1.03	0.21
(%)	(0.1)	(1.1)	(9.4)	(35.9)	(35.0)	(14.1)	(3.5)	(0.6)	(0.2)	(0.1)	(100.0)				
合計	363	8,392	50,797	70,263	28,134	4,901	552	79	25	38	163,544	9,445	172,989	0.86	0.18
(%)	(0.2)	(5.1)	(31.1)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	6	26	159	248	114	15	3				571	4,649	5,220	0.87	0.18
(%)	(1.1)	(4.6)	(27.8)	(43.4)	(20.0)	(2.6)	(0.5)				(100.0)				
総計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
平均	3.53	3.91	4.80	5.38	5.83	6.26	6.85	6.95	6.39	5.68	5.23	5.07	5.22		
標準偏差	1.75	1.24	1.23	1.30	1.46	1.72	2.12	2.50	2.49	2.26	1.42	1.50	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査