

表 3127 nPCR(g/kg/day) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前ヘモグロビン 濃度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
5未満	1	11	59	89	27	12	2				201	9	210	0.87	0.20
(%)	(0.5)	(5.5)	(29.4)	(44.3)	(13.4)	(6.0)	(1.0)				(100.0)				
5～	1	17	81	95	32	6					232	19	251	0.83	0.18
(%)	(0.4)	(7.3)	(34.9)	(40.9)	(13.8)	(2.6)					(100.0)				
6～	6	69	224	162	65	21	4		1		552	65	617	0.80	0.21
(%)	(1.1)	(12.5)	(40.6)	(29.3)	(11.8)	(3.8)	(0.7)		(0.2)		(100.0)				
7～	19	276	806	734	271	57	8	1		1	2,173	272	2,445	0.80	0.20
(%)	(0.9)	(12.7)	(37.1)	(33.8)	(12.5)	(2.6)	(0.4)	(0.0)		(0.0)	(100.0)				
8～	41	865	2,859	2,735	1,041	186	19	6	1	2	7,755	742	8,497	0.81	0.19
(%)	(0.5)	(11.2)	(36.9)	(35.3)	(13.4)	(2.4)	(0.2)	(0.1)	(0.0)	(0.0)	(100.0)				
9～	91	1,835	9,132	10,783	4,027	727	82	14	5	8	26,704	1,904	28,608	0.84	0.18
(%)	(0.3)	(6.9)	(34.2)	(40.4)	(15.1)	(2.7)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
10～	97	2,627	17,865	24,578	9,681	1,642	191	24	10	10	56,725	3,044	59,769	0.86	0.17
(%)	(0.2)	(4.6)	(31.5)	(43.3)	(17.1)	(2.9)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
11～	69	1,808	13,388	20,645	8,549	1,449	171	24	7	8	46,118	2,297	48,415	0.87	0.17
(%)	(0.1)	(3.9)	(29.0)	(44.8)	(18.5)	(3.1)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
12～	32	622	4,559	7,409	3,175	546	46	5		6	16,400	808	17,208	0.88	0.17
(%)	(0.2)	(3.8)	(27.8)	(45.2)	(19.4)	(3.3)	(0.3)	(0.0)		(0.0)	(100.0)				
13～	6	141	1,038	1,854	786	153	17	2	1	2	4,000	260	4,260	0.89	0.17
(%)	(0.2)	(3.5)	(26.0)	(46.4)	(19.7)	(3.8)	(0.4)	(0.1)	(0.0)	(0.1)	(100.0)				
14～	2	57	434	640	265	47	2				1,447	104	1,551	0.87	0.17
(%)	(0.1)	(3.9)	(30.0)	(44.2)	(18.3)	(3.2)	(0.1)				(100.0)				
合計	365	8,328	50,445	69,724	27,919	4,846	542	76	25	37	162,307	9,524	171,831	0.86	0.18
(%)	(0.2)	(5.1)	(31.1)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	4	90	511	787	329	70	13	3		1	1,808	4,570	6,378	0.88	0.19
(%)	(0.2)	(5.0)	(28.3)	(43.5)	(18.2)	(3.9)	(0.7)	(0.2)		(0.1)	(100.0)				
総計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
平均	10.14	10.32	10.62	10.79	10.84	10.82	10.74	10.66	10.36	10.73	10.72	10.51	10.71		
標準偏差	1.51	1.37	1.26	1.21	1.20	1.28	1.28	1.20	1.37	1.28	1.25	1.39	1.25		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査