

表 3130 nPCR(g/kg/day) Non-HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

Non-HDL-C濃度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
50未満	19	205	714	785	281	59	7	2	1	2	2,075	82	2,157	0.83	0.20
(%)	(0.9)	(9.9)	(34.4)	(37.8)	(13.5)	(2.8)	(0.3)	(0.1)	(0.0)	(0.1)	(100.0)				
50～	40	632	3,244	3,947	1,517	253	35	3		2	9,673	461	10,134	0.84	0.18
(%)	(0.4)	(6.5)	(33.5)	(40.8)	(15.7)	(2.6)	(0.4)	(0.0)		(0.0)	(100.0)				
70～	46	1,108	6,865	9,170	3,434	559	73	12	6	5	21,278	962	22,240	0.85	0.17
(%)	(0.2)	(5.2)	(32.3)	(43.1)	(16.1)	(2.6)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
90～	40	1,135	7,757	11,027	4,265	719	65	6	5	6	25,025	1,152	26,177	0.86	0.17
(%)	(0.2)	(4.5)	(31.0)	(44.1)	(17.0)	(2.9)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
110～	33	893	6,021	8,751	3,578	617	72	11	4	5	19,985	998	20,983	0.87	0.17
(%)	(0.2)	(4.5)	(30.1)	(43.8)	(17.9)	(3.1)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
130～	14	537	3,517	5,320	2,267	405	42	8	1	2	12,113	521	12,634	0.87	0.17
(%)	(0.1)	(4.4)	(29.0)	(43.9)	(18.7)	(3.3)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
150～	9	250	1,741	2,660	1,143	196	17	4	1	1	6,022	287	6,309	0.87	0.17
(%)	(0.1)	(4.2)	(28.9)	(44.2)	(19.0)	(3.3)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
170～	5	116	780	1,104	493	105	16		1		2,620	135	2,755	0.87	0.18
(%)	(0.2)	(4.4)	(29.8)	(42.1)	(18.8)	(4.0)	(0.6)		(0.0)		(100.0)				
190～	2	62	276	422	181	35	1				979	45	1,024	0.87	0.18
(%)	(0.2)	(6.3)	(28.2)	(43.1)	(18.5)	(3.6)	(0.1)				(100.0)				
210～	2	39	161	242	92	21	3				560	29	589	0.86	0.19
(%)	(0.4)	(7.0)	(28.8)	(43.2)	(16.4)	(3.8)	(0.5)				(100.0)				
合計	210	4,977	31,076	43,428	17,251	2,969	331	46	19	23	100,330	4,672	105,002	0.86	0.17
(%)	(0.2)	(5.0)	(31.0)	(43.3)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	159	3,441	19,880	27,083	10,997	1,947	224	33	6	15	63,785	9,422	73,207	0.86	0.18
(%)	(0.2)	(5.4)	(31.2)	(42.5)	(17.2)	(3.1)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
総計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
平均	94.36	102.55	104.80	106.84	108.34	109.55	107.47	105.76	101.84	97.13	106.31	106.98	106.34		
標準偏差	38.28	35.65	33.02	32.73	33.09	33.99	35.01	30.54	35.16	29.45	33.13	33.42	33.15		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査