

表 3133 nPCR(g/kg/day) 透析前脈拍(拍/分)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前脈拍	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
40未満	1	4	27	61	11	3			1		108	7	115	0.86	0.19
(%)	(0.9)	(3.7)	(25.0)	(56.5)	(10.2)	(2.8)			(0.9)		(100.0)				
40～	1	75	449	595	201	41	1	1		1	1,365	73	1,438	0.85	0.17
(%)	(0.1)	(5.5)	(32.9)	(43.6)	(14.7)	(3.0)	(0.1)	(0.1)		(0.1)	(100.0)				
50～	21	566	3,619	4,965	1,835	300	37	3		4	11,350	612	11,962	0.86	0.17
(%)	(0.2)	(5.0)	(31.9)	(43.7)	(16.2)	(2.6)	(0.3)	(0.0)		(0.0)	(100.0)				
60～	84	1,790	11,202	15,428	6,202	1,114	110	16	11	8	35,965	1,973	37,938	0.86	0.18
(%)	(0.2)	(5.0)	(31.1)	(42.9)	(17.2)	(3.1)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
70～	96	2,182	13,085	18,431	7,715	1,357	144	24	8	10	43,052	2,449	45,501	0.86	0.18
(%)	(0.2)	(5.1)	(30.4)	(42.8)	(17.9)	(3.2)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
80～	59	1,474	8,549	11,594	4,638	732	99	4	2	6	27,157	1,507	28,664	0.86	0.17
(%)	(0.2)	(5.4)	(31.5)	(42.7)	(17.1)	(2.7)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
90～	26	643	3,692	5,093	1,874	314	28	7		2	11,679	652	12,331	0.85	0.17
(%)	(0.2)	(5.5)	(31.6)	(43.6)	(16.0)	(2.7)	(0.2)	(0.1)		(0.0)	(100.0)				
100～	12	213	1,224	1,533	570	95	17			1	3,665	193	3,858	0.85	0.18
(%)	(0.3)	(5.8)	(33.4)	(41.8)	(15.6)	(2.6)	(0.5)			(0.0)	(100.0)				
110～	1	47	271	348	125	26	4	2			824	57	881	0.86	0.18
(%)	(0.1)	(5.7)	(32.9)	(42.2)	(15.2)	(3.2)	(0.5)	(0.2)			(100.0)				
120～	1	30	120	161	58	10	4			1	385	20	405	0.85	0.20
(%)	(0.3)	(7.8)	(31.2)	(41.8)	(15.1)	(2.6)	(1.0)			(0.3)	(100.0)				
合計	302	7,024	42,238	58,209	23,229	3,992	444	57	22	33	135,550	7,543	143,093	0.86	0.18
(%)	(0.2)	(5.2)	(31.2)	(42.9)	(17.1)	(2.9)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	67	1,394	8,718	12,302	5,019	924	111	22	3	5	28,565	6,551	35,116	0.86	0.18
(%)	(0.2)	(4.9)	(30.5)	(43.1)	(17.6)	(3.2)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
総計	369	8,418	50,956	70,511	28,248	4,916	555	79	25	38	164,115	14,094	178,209	0.86	0.18
(%)	(0.2)	(5.1)	(31.0)	(43.0)	(17.2)	(3.0)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
平均	74.95	75.12	74.63	74.50	74.47	74.17	75.67	74.09	68.50	74.15	74.56	74.65	74.56		
標準偏差	12.94	13.05	12.92	12.81	12.38	12.33	13.30	13.06	11.07	16.22	12.77	12.72	12.77		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査